

## mPRa Polyclonal Antibody

Catalog No: YT2825

**Reactivity:** Human; Mouse

**Applications:** WB;IHC

Target: mPRa

**Fields:** >>Chemical carcinogenesis - receptor activation

Gene Name: PAQR7

**Protein Name:** Membrane progestin receptor alpha

**Q86WK9** 

Q80ZE4

Human Gene Id: 164091

**Human Swiss Prot** 

idiliali Swiss Fiot

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

MPRA. AA range:297-346

**Specificity:** mPRa Polyclonal Antibody detects endogenous levels of mPRa protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000;IHC 1:50-300

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 40kD

**Background:** function: Steroid membrane receptor. Binds progesterone in vitro. May be

involved in oocyte maturation., similarity: Belongs to the ADIPOR family., tissue

specificity:Expressed in ovary, testis, placenta, uterus and bladder.,

**Function:** function: Steroid membrane receptor. Binds progesterone in vitro. May be

involved in oocyte maturation., similarity: Belongs to the ADIPOR family., tissue

specificity: Expressed in ovary, testis, placenta, uterus and bladder.,


Subcellular Location:

Cell membrane; Multi-pass membrane protein.


**Expression:** Expressed in a wide range of tissues including ovary, testis, placenta, uterus

and bladder.


## **Products Images**



Western Blot analysis of 293 cells using mPRa Polyclonal Antibody



Western blot analysis of lysates from 293 and COLO cells, using MPRA Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).