

NCoA-5 Polyclonal Antibody

Catalog No :	YT2997
Reactivity :	Human;Mouse
Applications :	IHC;IF;ELISA
Target :	NCoA-5
Gene Name :	NCOA5
Protein Name :	Nuclear receptor coactivator 5
Human Gene Id :	57727
Human Swiss Prot No :	Q9HCD5
Mouse Gene Id :	228869
Mouse Swiss Prot No :	Q91W39
Immunogen :	The antiserum was produced against synthesized peptide derived from human NCOA5. AA range:345-394
Specificity :	NCoA-5 Polyclonal Antibody detects endogenous levels of NCoA-5 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 65kD

Background :

This gene encodes a coregulator for the alpha and beta estrogen receptors and the orphan nuclear receptor NR1D2. The protein localizes to the nucleus, and is thought to have both coactivator and corepressor functions. Its interaction with nuclear receptors is independent of the AF2 domain on the receptors, which is known to regulate interaction with other coreceptors. Two alternatively spliced transcript variants for this gene have been described. However, the full length nature of one of the variants has not been determined. [provided by RefSeq, Jul 2008],

Function :

domain:Contains one Leu-Xaa-Xaa-Leu-Leu (LxxLL) motif that is essential for the association with nuclear receptors.,function:Nuclear receptor coregulator that can have both coactivator and corepressor functions. Interacts with nuclear receptors for steroids (ESR1 and ESR2) independently of the steroid binding domain (AF-2) of the ESR receptors, and with the orphan nuclear receptor NR1D2. Involved in the coactivation of nuclear steroid receptors (ER) as well as the corepression of MYC/c-myc in response to 17-beta-estradiol (E2).,subunit:Binds HTATIP2/TIP30.,tissue specificity:Widely expressed.,


Subcellular Location :

Nucleus.

Expression :

Widely expressed.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using NCOA5 Antibody. The picture on the right is blocked with the synthesized peptide.