

Neurexin III β Polyclonal Antibody

Catalog No :	YT3057
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA;IHC
Target :	Neurexin III β
Gene Name :	NRXN3
Protein Name :	Neurexin-3-beta
Human Gene Id :	9369
Human Swiss Prot No :	Q9HDB5
Mouse Swiss Prot No :	Q8C985
Immunogen :	Synthesized peptide derived from Neurexin III β . at AA range: 30-110
Specificity :	Neurexin III β Polyclonal Antibody detects endogenous levels of Neurexin III β protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	70kD

Cell Pathway : Cell adhesion molecules (CAMs);


Background : NRXN3 (neurexin 3) encodes a member of a family of proteins that function in the nervous system as receptors and cell adhesion molecules. Extensive alternative splicing and the use of alternative promoters results in multiple transcript variants and protein isoforms for this gene, but the full-length nature of many of these variants has not been determined. Transcripts that initiate from an upstream promoter encode alpha isoforms, which contain epidermal growth factor-like (EGF-like) sequences and laminin G domains. Transcripts initiating from the downstream promoter encode beta isoforms, which lack EGF-like sequences. Genetic variation at this locus has been associated with a range of behavioral phenotypes, including alcohol dependence and autism spectrum disorder.


Function : cell morphogenesis, cell morphogenesis involved in differentiation, regulation of neurotransmitter levels, generation of a signal involved in cell-cell signaling, neurotransmitter transport, cell motion, cell adhesion, cell-cell signaling, synaptic transmission, neurotransmitter secretion, axonogenesis, axon guidance, synaptogenesis, transmission of nerve impulse, biological adhesion, cell projection organization, neuron differentiation, neuron projection development, secretion by cell, cellular component morphogenesis, cell part morphogenesis, extracellular structure organization, secretion, neuron development, cell morphogenesis involved in neuron differentiation, neuron projection morphogenesis, cell projection morphogenesis, synapse organization, neurological system process,

Subcellular Location : Membrane ; Single-pass type I membrane protein .

Expression : Expressed in the blood vessel walls (at protein level).

Products Images





Immunohistochemical analysis of paraffin-embedded human lung cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).