

Olfactory receptor 52A5 Polyclonal Antibody

YT3375 Catalog No:

Reactivity: Human

WB;IF;ELISA **Applications:**

Olfactory receptor 52A5 **Target:**

Fields: >>Olfactory transduction

Gene Name: OR52A5

Protein Name: Olfactory receptor 52A5

Q9H2C5

Human Gene Id: 390054

Human Swiss Prot

No:

The antiserum was produced against synthesized peptide derived from human Immunogen:

OR52A5. AA range:160-209

Olfactory receptor 52A5 Polyclonal Antibody detects endogenous levels of **Specificity:**

Olfactory receptor 52A5 protein.

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Formulation:

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/2



Observed Band: 36kD

Cell Pathway: Olfactory transduction;

Background:

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],


Function:

function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family.,


Subcellular Location:

Cell membrane; Multi-pass membrane protein.

Products Images



Western blot analysis of lysates from Jurkat cells, using OR52A5 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using OR52A5 antibody.