

ORC1 Polyclonal Antibody

Catalog No :	YT3469
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	ORC1
Fields :	>>Cell cycle
Gene Name :	ORC1
Protein Name :	Origin recognition complex subunit 1
Human Gene Id :	4998
Human Swiss Prot No :	Q13415
Mouse Swiss Prot No :	Q9Z1N2
Immunogen :	The antiserum was produced against synthesized peptide derived from human ORC1L. AA range:331-380
Specificity :	ORC1 Polyclonal Antibody detects endogenous levels of ORC1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 100,120kD

Cell Pathway : Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;


Background : The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is the largest subunit of the ORC complex. While other ORC subunits are stable throughout the cell cycle, the levels of this protein vary during the cell cycle, which has been shown to be controlled by ubiquitin-mediated proteolysis after initiation of DNA replication. This protein is found to be selectively phosphorylated during mitosis. It is also reported to interact with MYST histone acetyltransferase 2 (MyST2/HBO1), a protein involved in control of transcription silencing. Alternatively spliced transcr

Function : function:Component of the origin recognition complex (ORC) that binds origins of replication. It has a role in both chromosomal replication and mating type transcriptional silencing. Binds to the ARS consensus sequence (ACS) of origins of replication in an ATP-dependent manner.,similarity:Belongs to the ORC1 family.,similarity:Contains 1 BAH domain.,subunit:ORC is composed of six subunits. Interacts with CDC6 and MYST2/HBO1.,


Subcellular Location : Nucleus.

Expression : Epithelium, Eye,


Products Images



Western Blot analysis of various cells using ORC1 Polyclonal Antibody



Western Blot analysis of 3T3 cells using ORC1 Polyclonal Antibody



Western blot analysis of the lysates from HepG2 cells using ORC1L antibody.