

## Pax-9 Polyclonal Antibody

Catalog No: YT3603

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA

Target: Pax-9

Gene Name: PAX9

**Protein Name:** Paired box protein Pax-9

P55771

P47242

Human Gene Id: 5083

**Human Swiss Prot** 

No:

Mouse Gene ld: 18511

**Mouse Swiss Prot** 

No:

**Rat Gene Id:** 362741

Rat Swiss Prot No: Q2L4T2

Immunogen: The antiserum was produced against synthesized peptide derived from human

Pax-9. AA range:158-207

**Specificity:** Pax-9 Polyclonal Antibody detects endogenous levels of Pax-9 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



**Concentration**: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 35kD

**Background:** This gene is a member of the paired box (PAX) family of transcription factors.

Members of this gene family typically contain a paired box domain, an octapeptide, and a paired-type homeodomain. These genes play critical roles during fetal development and cancer growth. Mice lacking this gene exhibit impaired development of organs, musculature and the skeleton, including absent and abnormally developed teeth, and neonatal lethality. Mutations in the human gene are associated with selective tooth agenesis-3. [provided by RefSeq, Sep

2015],

**Function:** disease:Defects in PAX9 are a cause of oligodontia [MIM:604625]. It is a form of

familial or selective tooth agenesis. Oligodontia is defined as the agenesis of 6 or more permanent teeth without associated systemic disorders. Agenesis of one or more teeth constitutes one of the most common developmental anomalies in man. Reported incidences vary from 1.6% to 9.6%, excluding third molar (Wisdom tooth) agenesis, which occurs in 20% of the population., function: Transcription

factor required for normal development of thymus, parathyroid glands,


ultimobranchial bodies, teeth, skeletal elements of skull and larynx as well as distal limbs.,similarity:Contains 1 paired domain.,subunit:Interacts with KDM5B.,

Subcellular Location:

Nucleus.

**Expression:** Lung, Oesophagus,

## **Products Images**



Western Blot analysis of various cells using Pax-9 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



