


PLC β 3 Polyclonal Antibody


Catalog No :	YT3790
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PLC β 3
Fields :	>>Inositol phosphate metabolism;>>Metabolic pathways;>>Rap1 signaling pathway;>>Calcium signaling pathway;>>cGMP-PKG signaling pathway;>>Chemokine signaling pathway;>>Phosphatidylinositol signaling system;>>Sphingolipid signaling pathway;>>Phospholipase D signaling pathway;>>Adrenergic signaling in cardiomyocytes;>>Vascular smooth muscle contraction;>>Wnt signaling pathway;>>Apelin signaling pathway;>>Gap junction;>>Platelet activation;>>Neutrophil extracellular trap formation;>>NOD-like receptor signaling pathway;>>Circadian entrainment;>>Long-term potentiation;>>Retrograde endocannabinoid signaling;>>Glutamatergic synapse;>>Cholinergic synapse;>>Serotonergic synapse;>>Dopaminergic synapse;>>Long-term depression;>>Taste transduction;>>Inflammatory mediator regulation of TRP channels;>>Insulin secretion;>>GnRH signaling pathway;>>Estrogen signaling pathway;>>Melanogenesis;>>Thyroid hormone synthesis;>>Thyroid hormone signaling pathway;>>Oxytocin signaling pathway;>>Glucagon signaling p
Gene Name :	PLCB3
Protein Name :	1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase beta-3
Human Gene Id :	5331
Human Swiss Prot No :	Q01970
Mouse Swiss Prot No :	P51432
Rat Swiss Prot No :	Q99JE6
Immunogen :	The antiserum was produced against synthesized peptide derived from human PLC beta3. AA range:503-552
Specificity :	PLC β 3 Polyclonal Antibody detects endogenous levels of PLC β 3 protein.

Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	140kD
Cell Pathway :	Stem cell pathway; WNT;WNT-T CELL;β-Catenin; AMPK
Background :	This gene encodes a member of the phosphoinositide phospholipase C beta enzyme family that catalyze the production of the secondary messengers diacylglycerol and inositol 1,4,5-triphosphate from phosphatidylinositol in G-protein-linked receptor-mediated signal transduction. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010],
Function :	catalytic activity:1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate + H(2)O = 1D-myo-inositol 1,4,5-trisphosphate + diacylglycerol.,cofactor:Calcium.,function:The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes.,similarity:Contains 1 C2 domain.,similarity:Contains 1 PI-PLC X-box domain.,similarity:Contains 1 PI-PLC Y-box domain.,subunit:Interacts with SHANK2 (By similarity). Interacts with LPAR2.,
Subcellular Location :	Cytoplasm . Membrane . Nucleus . And particulate fractions. .
Expression :	Epithelium,Uterus,

Products Images



Western Blot analysis of various cells using PLC β 3 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PLC beta3 Antibody. The picture on the right is blocked with the synthesized peptide.