

PDK1 Polyclonal Antibody

Catalog No: YT3644

Reactivity: Human; Mouse; Rat

Applications: WB;IHC

Target: PDK1

Fields: >>HIF-1 signaling pathway;>>Central carbon metabolism in cancer

Gene Name: PDK1

Protein Name: [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 1 mitochondrial

Human Gene Id: 5163

Human Swiss Prot

No:

Mouse Gene ld: 228026

Mouse Swiss Prot

No:

Rat Swiss Prot No: Q63065

Immunogen: Synthesized peptide derived from PDK1 . at AA range: 80-160

Specificity: PDK1 Polyclonal Antibody detects endogenous levels of PDK1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500-2000;IHC 1:50-300

Q15118

Q8BFP9

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 50kD

Cell Pathway : T_Cell_Receptor;Fc epsilon RI;Neurotrophin;

Background: Pyruvate dehydrogenase (PDH) is a mitochondrial multienzyme complex that

catalyzes the oxidative decarboxylation of pyruvate and is one of the major enzymes responsible for the regulation of homeostasis of carbohydrate fuels in

mammals. The enzymatic activity is regulated by a

phosphorylation/dephosphorylation cycle. Phosphorylation of PDH by a specific

pyruvate dehydrogenase kinase (PDK) results in inactivation. Multiple

alternatively spliced transcript variants have been found for this gene. [provided

by RefSeq, Jun 2013],

Function: catalytic activity:ATP + [pyruvate dehydrogenase (acetyl-transferring)] = ADP +

[pyruvate dehydrogenase (acetyl-transferring)] phosphate.,function:Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1

alpha subunit, thus contributing to the regulation of glucose

metabolism.,similarity:Belongs to the PDK/BCKDK protein kinase family.,similarity:Contains 1 histidine kinase domain.,tissue specificity:Expressed

predominantly in the heart.,


Subcellular Location:

Mitochondrion matrix.


Expression: Expressed predominantly in the heart. Detected at lower levels in liver, skeletal

muscle and pancreas.

Products Images Western Blot analysis of various cells using PDK1 Polyclonal Antibody -PDK1 39



Western Blot analysis of Hela cells using PDK1 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at $1:200(4^{\circ}$ overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min).