

## PKR Polyclonal Antibody

Catalog No: YT3779

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;IHC;IF;ELISA

Target: PKR

Fields: >>Protein processing in endoplasmic reticulum;>>Necroptosis;>>Alzheimer

disease;>>Hepatitis C;>>Measles;>>Influenza A;>>Human papillomavirus infection;>>Kaposi sarcoma-associated herpesvirus infection;>>Herpes simplex

virus 1 infection;>>Epstein-Barr virus infection;>>Coronavirus disease -

COVID-19;>>Viral carcinogenesis

Gene Name: EIF2AK2

**Protein Name:** Interferon-induced double-stranded RNA-activated protein kinase

Human Gene Id: 5610

**Human Swiss Prot** 

No:

Mouse Swiss Prot

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

PKR. AA range:413-462

P19525

Q03963

**Specificity:** PKR Polyclonal Antibody detects endogenous levels of PKR protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 74kD

**Background:** The protein encoded by this gene is a serine/threonine protein kinase that is

activated by autophosphorylation after binding to dsRNA. The activated form of the encoded protein can phosphorylate translation initiation factor EIF2S1, which in turn inhibits protein synthesis. This protein is also activated by manganese ions and heparin. Three transcript variants encoding two different isoforms have been

found for this gene. [provided by RefSeq, Oct 2011],

**Function:** catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme

regulation: Activity is markedly stimulated by manganese ions. Besides dsRNA, heparin is a potent activator of the kinase. Binding to dsRNA is required for dimerization leading to autophosphorylation in the activation loop and stimulation

of function. Inhibited by vaccinia virus protein E3, probably via dsRNA sequestering.,function:Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated

during the course of a viral infection.,induction:By

interferon.,PTM:Autophosphorylated on several Ser and Thr residues.

Autophosphorylation of Thr-451 is dependent on Thr-446 and is stimulated by

dsRNA bindin

Subcellular Location : Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Nuclear localization is elevated in acute leukemia, myelodysplastic syndrome (MDS), melanoma, breast, colon, prostate and lung cancer patient samples or cell lines as well as neurocytes


from advanced Creutzfeldt-Jakob disease patients...

**Expression:** Highly expressed in thymus, spleen and bone marrow compared to non-


hematopoietic tissues such as small intestine, liver, or kidney tissues. Colocalizes with GSK3B and TAU in the Alzheimer disease (AD) brain. Elevated levels seen in breast and colon carcinomas, and which correlates with tumor progression and

invasiveness or risk of progression.


## **Products Images**




Western Blot analysis of various cells using PKR Polyclonal Antibody diluted at 1:2000




Western Blot analysis of HeLa cells using PKR Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PKR Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa, using PKR Antibody. The lane on the right is blocked with the synthesized peptide.



Feng, Qin. "Quantitative proteomic analysis reveals that Arctigenin alleviates concanavalin A-induced hepatitis through suppressing immune system and regulating autophagy." Frontiers in immunology 9 (2018): 1881.