


WNT8A Polyclonal Antibody

Catalog No :	YN0289
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	WNT8A
Fields :	>>mTOR signaling pathway;>>Wnt signaling pathway;>>Hippo signaling pathway;>>Signaling pathways regulating pluripotency of stem cells;>>Melanogenesis;>>Cushing syndrome;>>Alzheimer disease;>>Pathways of neurodegeneration - multiple diseases;>>Human papillomavirus infection;>>Pathways in cancer;>>Proteoglycans in cancer;>>Basal cell carcinoma;>>Breast cancer;>>Hepatocellular carcinoma;>>Gastric cancer
Gene Name :	WNT8A WNT8D
Protein Name :	Protein Wnt-8a (Protein Wnt-8d)
Human Gene Id :	7478
Human Swiss Prot	Q9H1J5
No : Mouse Swiss Prot	Q64527
No : Immunogen :	Synthesized peptide derived from human protein . at AA range: 90-170
Specificity :	WNT8A Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.



Best Tools for immunology Research	
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	38kD
Cell Pathway :	WNT;WNT-T CELLHedgehog;Melanogenesis;Pathways in cancer;Basal cell carcinoma;
Background :	The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family, and may be implicated in development of early embryos as well as germ cell tumors. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2014],
Function :	function:Ligand for members of the frizzled family of seven transmembrane receptors. May play an important role in the development and differentiation of certain forebrain structures, notably the hippocampus.,similarity:Belongs to the Wnt family.,
Subcellular Location :	Secreted, extracellular space, extracellular matrix . Secreted .



Products Images

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night