

**Ribosomal Protein L7L1 Polyclonal Antibody**


|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YT4119  |
| <b>Reactivity :</b>          | Human;Mouse   |
| <b>Applications :</b>        | WB;IHC;IF;ELISA   |
| <b>Target :</b>              | Ribosomal Protein L7L1  |
| <b>Gene Name :</b>           | RPL7L1  |
| <b>Protein Name :</b>        | 60S ribosomal protein L7-like 1   |
| <b>Human Gene Id :</b>       | 285855  |
| <b>Human Swiss Prot No :</b> | Q6DKI1  |
| <b>Mouse Swiss Prot No :</b> | Q9D8M4  |
| <b>Immunogen :</b>           | The antiserum was produced against synthesized peptide derived from human RPL7L1. AA range:41-90                      |
| <b>Specificity :</b>         | Ribosomal Protein L7L1 Polyclonal Antibody detects endogenous levels of Ribosomal Protein L7L1 protein.               |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200  |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Concentration :</b>       | 1 mg/ml   |
| <b>Storage Stability :</b>   | -15°C to -25°C/1 year(Do not lower than -25°C)  |
| <b>Observed Band :</b>       | 29kD  |

---


|                               |  |
|-------------------------------|--|
| <b>Background :</b>           | <u>similarity:Belongs to the ribosomal protein L30P family.,</u> |
| <b>Function :</b>             | <u>similarity:Belongs to the ribosomal protein L30P family.,</u> |
| <b>Subcellular Location :</b> | <u>nucleolus,ribosome,cytosolic large ribosomal subunit,</u>     |
| <b>Expression :</b>           | <u>Eye,</u>  |

---

## Products Images



Western blot analysis of lysates from HepG2 cells, using RPL7L1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).