

Sp2 Polyclonal Antibody

Catalog No :	YT4375
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	Sp2
Gene Name :	SP2
Protein Name :	Transcription factor Sp2
Human Gene Id :	6668
Human Swiss Prot No :	Q02086
Mouse Gene Id :	78912
Mouse Swiss Prot No :	Q9D2H6
Immunogen :	The antiserum was produced against synthesized peptide derived from human SP2. AA range:471-520
Specificity :	Sp2 Polyclonal Antibody detects endogenous levels of Sp2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 64kD

Background :

This gene encodes a member of the Sp subfamily of Sp/XKLF transcription factors. Sp family proteins are sequence-specific DNA-binding proteins characterized by an amino-terminal trans-activation domain and three carboxy-terminal zinc finger motifs. This protein contains the least conserved DNA-binding domain within the Sp subfamily of proteins, and its DNA sequence specificity differs from the other Sp proteins. It localizes primarily within subnuclear foci associated with the nuclear matrix, and can activate or in some cases repress expression from different promoters. [provided by RefSeq, Jul 2008],

Function :

function: Binds to GC box promoters elements and selectively activates mRNA synthesis from genes that contain functional recognition sites., similarity: Belongs to the Sp1 C2H2-type zinc-finger protein family., similarity: Contains 3 C2H2-type zinc fingers.,



Subcellular Location :


Nucleus.

Expression :

Bone marrow, Brain, Lung, Lymph, Spleen,

Products Images





Immunohistochemical analysis of paraffin-embedded human spleen. 1, Tris-EDTA, pH 9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).