

StARD13 Polyclonal Antibody

| | |
|------------------------------|---|
| Catalog No : | YT4438 |
| Reactivity : | Human;Mouse |
| Applications : | WB;ELISA;IHC |
| Target : | StARD13 |
| Gene Name : | STARD13 |
| Protein Name : | StAR-related lipid transfer protein 13 |
| Human Gene Id : | 90627 |
| Human Swiss Prot No : | Q9Y3M8 |
| Mouse Gene Id : | 243362 |
| Mouse Swiss Prot No : | Q923Q2 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human STA13. AA range:101-150 |
| Specificity : | StARD13 Polyclonal Antibody detects endogenous levels of StARD13 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |

Observed Band : 125kD


Background : This gene encodes a protein which contains an N-terminal sterile alpha motif (SAM) for protein-protein interactions, followed by an ATP/GTP-binding motif, a GTPase-activating protein (GAP) domain, and a C-terminal STAR-related lipid transfer (START) domain. It may be involved in regulation of cytoskeletal reorganization, cell proliferation, and cell motility, and acts as a tumor suppressor in hepatoma cells. The gene is located in a region of chromosome 13 that is associated with loss of heterozygosity in hepatocellular carcinomas. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Aug 2011],

Function : function:GTPase-activating protein for RhoA, and perhaps for Cdc42. May be involved in regulation of cytoskeletal reorganization, cell proliferation and cell motility. Acts a tumor suppressor in hepatocellular carcinoma cells.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 1 START domain.,subunit:Homodimer. Interacts with TAX1BP1.,tissue specificity:Ubiquitously expressed. Underexpressed in hepatocellular carcinoma cells and some breast cancer cell lines.,


Subcellular Location : Cytoplasm. Membrane; Peripheral membrane protein; Cytoplasmic side. Mitochondrion membrane; Peripheral membrane protein; Cytoplasmic side. Lipid droplet.

Expression : Ubiquitously expressed. Underexpressed in hepatocellular carcinoma cells and some breast cancer cell lines.


Products Images



Western blot analysis of STA13 Antibody. The lane on the right is blocked with the STA13 peptide.



Immunofluorescence analysis of A549 cells, using STA13 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).