

**PI 3 Kinase P85α Monoclonal Antibody(3B7)**

<b>Catalog No :</b>	YM3503
<b>Reactivity :</b>	Rat;Mouse
<b>Applications :</b>	WB;IHC;IF
<b>Target :</b>	PI3 Kinase P85α
<b>Fields :</b>	>>EGFR tyrosine kinase inhibitor resistance;>>Endocrine resistance;>>Platinum drug resistance;>>ErbB signaling pathway;>>Ras signaling pathway;>>Rap1 signaling pathway;>>cAMP signaling pathway;>>Chemokine signaling pathway;>>HIF-1 signaling pathway;>>FoxO signaling pathway;>>Phosphatidylinositol signaling system;>>Sphingolipid signaling pathway;>>Phospholipase D signaling pathway;>>Autophagy - animal;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>AMPK signaling pathway;>>Apoptosis;>>Longevity regulating pathway;>>Longevity regulating pathway - multiple species;>>Cellular senescence;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast differentiation;>>Focal adhesion;>>Signaling pathways regulating pluripotency of stem cells;>>Platelet activation;>>Neutrophil extracellular trap formation;>>Toll-like receptor signaling pathway;>>C-type lectin receptor signaling pathway;>>JAK-STAT signaling pathway;>>Natural killer cell mediated cytotoxicity;>>T cell receptor signaling pathway;>
<b>Gene Name :</b>	PIK3R1
<b>Protein Name :</b>	Phosphatidylinositol 3-kinase regulatory subunit alpha (PI3-kinase regulatory subunit alpha) (PI3K regulatory subunit alpha) (PtdIns-3-kinase regulatory subunit alpha) (Phosphatidylinositol 3-kinase 8
<b>Human Gene Id :</b>	5295
<b>Human Swiss Prot No :</b>	P27986
<b>Mouse Swiss Prot No :</b>	P26450
<b>Rat Swiss Prot No :</b>	Q63787
<b>Immunogen :</b>	Recombinant Protein of PI3 Kinase P85α


---

<b>Specificity :</b>	<u>PI3 Kinase P85<math>\alpha</math> protein detects endogenous levels of PI3 Kinase P85<math>\alpha</math></u>
<b>Formulation :</b>	<u>Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.</u>
<b>Source :</b>	<u>Monoclonal, Mouse</u>
<b>Dilution :</b>	<u>WB 1:1000-2000, IHC 1:100-200. IF 1:50-200</u>
<b>Purification :</b>	<u>The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.</u>
<b>Concentration :</b>	<u>1 mg/ml</u>
<b>Storage Stability :</b>	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
<b>Observed Band :</b>	<u>85kD</u>
<b>Cell Pathway :</b>	<u>ErbB_HER;Chemokine;Phosphatidylinositol signaling system;mTOR;Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;VEGF;Focal adhesion;Toll_Like;Jak_STAT;Natural killer cell mediated cyto</u>
<b>Background :</b>	<u>Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different isoforms. [provided by RefSeq, Jun 2011],</u>
<b>Function :</b>	<u>disease:Defects in PIK3R1 are a cause of severe insulin resistance.,domain:The SH3 domain mediates the binding to CBLB, and to HIV-1 Nef.,function:Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues.,PTM:Polyubiquitinated in T-cells by CBLB; which does not promote proteasomal degradation but impairs association with CD28 and CD3Z upon T-cell activation.,similarity:Belongs to the PI3K p85 subunit family.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SH2 domains.,subunit:Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunits. Interacts with phosphorylated TOM1L1. Interacts with phosphorylat</u>
<b>Subcellular Location :</b>	<u>nucleus,cytoplasm,cis-Golgi network,cytosol,plasma membrane,cell-cell junction,phosphatidylinositol 3-kinase complex,phosphatidylinositol 3-kinase complex, class IA,membrane,perinuclear endoplasmic reticulum membrane,</u>


---

**Expression :** Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level).


## Products Images




Wang, L., Xiong, Q., Li, P. et al. The negative charge of the 343 site is essential for maintaining physiological functions of CXCR4. *BMC Mol and Cell Biol* 22, 8 (2021).



Western blot analysis of 1) 3T3, 2) Rat Liver Tissue with PI3 Kinase P85α Mouse mAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Rat Liver Tissue using PI3 Kinase P85α Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue using PI3 Kinase P85 α Mouse mAb diluted at 1:200.