

## v-SNARE Vti1a Polyclonal Antibody

Catalog No: YT4894

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA

Target: v-SNARE Vti1a

**Fields:** >>SNARE interactions in vesicular transport

Q96AJ9

089116

Gene Name: VTI1A

Protein Name: Vesicle transport through interaction with t-SNAREs homolog 1A

Human Gene Id: 143187

**Human Swiss Prot** 

No:

Mouse Gene ld: 53611

**Mouse Swiss Prot** 

No:

Rat Gene ld: 65277

Rat Swiss Prot No: Q9JI51

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

VTI1A. AA range:31-80

Specificity: v-SNARE Vti1a Polyclonal Antibody detects endogenous levels of v-SNARE

Vti1a protein.

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.

1/3



**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 23kD

**Cell Pathway:** SNARE interactions in vesicular transport;

**Background:** The protein encoded by this gene is a member of the family of soluble N-

ethylmaleimide-sensitive fusion protein-attachment protein receptors (SNAREs) that function in intracellular trafficking. This family member is involved in vesicular transport between endosomes and the trans-Golgi network. It is a vesicle-associated SNARE (v-SNARE) that interacts with target membrane SNAREs (t-

SNAREs). Polymorphisms in this gene have been associated with binocular function, and also with susceptibility to colorectal and lung cancers. A recurrent rearrangement has been found between this gene and the transcription factor 7-like 2 (TCF7L2) gene in colorectal cancers. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Dec 2015],

**Function:** function:V-SNARE that mediates vesicle transport pathways through

interactions with t-SNAREs on the target membrane. These interactions are proposed to mediate aspects of the specificity of vesicle trafficking and to promote fusion of the lipid bilayers. May be concerned with increased secretion of cytokines associated with cellular senescence..similarity:Belongs to the VTI1

family.,


Subcellular Cytoplasmic vesicle . Golgi apparatus membrane ; Single-pass type IV

**Location:** membrane protein.


**Expression:** Lung,

## **Products Images**

2/3



Western blot analysis of VTI1A Antibody. The lane on the right is blocked with the VTI1A peptide.



Western blot analysis of the lysates from HUVECcells using VTI1A antibody.