


UDP-GlcDH Polyclonal Antibody


| | |
|------------------------------|---|
| Catalog No : | YT4816 |
| Reactivity : | Human;Mouse;Rat |
| Applications : | WB;IHC;IF;ELISA |
| Target : | UDP-GlcDH |
| Fields : | >>Pentose and glucuronate interconversions;>>Ascorbate and aldarate metabolism;>>Amino sugar and nucleotide sugar metabolism;>>Metabolic pathways;>>Biosynthesis of cofactors;>>Biosynthesis of nucleotide sugars |
| Gene Name : | UGDH |
| Protein Name : | UDP-glucose 6-dehydrogenase |
| Human Gene Id : | 7358 |
| Human Swiss Prot No : | O60701 |
| Mouse Gene Id : | 22235 |
| Mouse Swiss Prot No : | O70475 |
| Rat Gene Id : | 83472 |
| Rat Swiss Prot No : | O70199 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human UGDH. AA range:391-440 |
| Specificity : | UDP-GlcDH Polyclonal Antibody detects endogenous levels of UDP-GlcDH protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |

| | |
|-------------------------------|---|
| Dilution : | WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15 °C to -25 °C/1 year(Do not lower than -25 °C) |
| Observed Band : | 55kD |
| Cell Pathway : | Pentose and glucuronate interconversions;Ascorbate and aldarate metabolism;Starch and sucrose metabolism;Amino sugar and nucleotide sugar metabolism; |
| Background : | The protein encoded by this gene converts UDP-glucose to UDP-glucuronate and thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin sulfate, and heparan sulfate. These glycosylated compounds are common components of the extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth and metastasis. The expression of this gene is up-regulated by transforming growth factor beta and down-regulated by hypoxia. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010], |
| Function : | catalytic activity:UDP-glucose + 2 NAD(+) + H(2)O = UDP-glucuronate + 2 NADH.,function:Involved in the biosynthesis of glycosaminoglycans; hyaluronan, chondroitin sulfate, and heparan sulfate.,pathway:Nucleotide-sugar biosynthesis; UDP-glucuronic acid biosynthesis; UDP-glucuronic acid from UDP-glucose: step 1/1.,similarity:Belongs to the UDP-glucose/GDP-mannose dehydrogenase family.,subunit:Homohexamer., |
| Subcellular Location : | nucleus,nucleoplasm,cytosol,extracellular exosome, |
| Expression : | Detected in heart, placenta, liver, pancreas, spleen, thymus, prostate, ovary, small intestine and colon (PubMed:9737970). Widely expressed (PubMed:9737970). |

Products Images



Western blot analysis of lysates from COLO cells, using UGDH Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).