

CD177 Polyclonal Antibody

Catalog No :	YT5283
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	CD177
Gene Name :	CD177
Protein Name :	CD177 antigen
Human Gene Id :	57126
Human Swiss Prot No :	Q8N6Q3
Mouse Gene Id :	68891
Mouse Swiss Prot No :	Q8R2S8
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human CD177. AA range:361-410
Specificity :	CD177 Polyclonal Antibody detects endogenous levels of CD177 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 46kD

Background : This gene encodes a glycosyl-phosphatidylinositol (GPI)-linked cell surface glycoprotein that plays a role in neutrophil activation. The protein can bind platelet endothelial cell adhesion molecule-1 and function in neutrophil transmigration. Mutations in this gene are associated with myeloproliferative diseases. Over-expression of this gene has been found in patients with polycythemia rubra vera. Autoantibodies against the protein may result in pulmonary transfusion reactions, and it may be involved in Wegener's granulomatosis. A related pseudogene, which is adjacent to this gene on chromosome 19, has been identified. [provided by RefSeq, Apr 2014],

Function : induction:By CSF3 in resting granulocytes. Induced in patients with polycythemia vera (PV) and with essential thrombocythemia (ET).,mass spectrometry: PubMed:11465086,polymorphism:There is a significant association between the variants Ala-3, Leu-251 and Thr-348 and a low expression of CD177 on neutrophils.,PTM:A soluble form may also be produced by proteolytic cleavage at the cell surface (shedding).,PTM:N-glycosylated.,similarity:Contains 2 UPAR/Ly6 domains.,tissue specificity:Highly expressed in normal bone marrow and weakly expressed in fetal liver. Expressed on neutrophils. Expressed in granulocytes of patients with polycythemia vera (PV) and with essential thrombocythemia (ET).,


Subcellular Location : Cell membrane ; Lipid-anchor, GPI-anchor . Membrane raft ; Lipid-anchor, GPI-like-anchor . Secreted . Cytoplasmic granule membrane . Cell projection, lamellipodium . Cell surface expression on neutrophils is increased upon TNF-alpha, fMLP or CXCL8/IL8-mediated stimulation (PubMed:17244676, PubMed:17580308). In neutrophils, stored predominantly in secondary and tertiary granules (PubMed:18462208). Can also be shedded from the cell membrane (PubMed:12239154, PubMed:18462208). Localizes to lamellar protrusions in spreading neutrophils (PubMed:28807980). .

Expression : Highly expressed in normal bone marrow and weakly expressed in fetal liver (PubMed:10753836). During neutrophil differentiation, expression begins at the metamyelocyte stage and continues throughout the subsequent stages (at protein level) (PubMed:17244676, PubMed:18462208, PubMed:24926686). Expressed by a subset of mature neutrophils (at protein level) (PubMed:10753836, PubMed:28240246, PubMed:12377969, PubMed:18462208, PubMed:12675722, PubMed:17244676, PubMed:17580308, PubMed:21193407, PubMed:24926686, PubMed:28807980, PubMed:27227454). The percentage of neutrophils expressing CD177 varies across the population (PubMed:17244676, PubMed:27227454). Expressed in granulocytes of patients with polycythemia vera (PV) and with essential thrombocythemia (ET) (PubMed:10753836, PubMed:12377969).


Products Images



Western Blot analysis of K562 cells using CD177 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



Western blot analysis of lysate from K562 cells, using CD177 Antibody.