


CD298 Polyclonal Antibody

Catalog No :	YT5623
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	CD298
Fields :	>>cGMP-PKG signaling pathway;>>cAMP signaling pathway;>>Cardiac muscle contraction;>>Adrenergic signaling in cardiomyocytes;>>Insulin secretion;>>Thyroid hormone synthesis;>>Thyroid hormone signaling pathway;>>Aldosterone synthesis and secretion;>>Aldosterone-regulated sodium reabsorption;>>Endocrine and other factor-regulated calcium reabsorption;>>Proximal tubule bicarbonate reclamation;>>Salivary secretion;>>Gastric acid secretion;>>Pancreatic secretion;>>Carbohydrate digestion and absorption;>>Protein digestion and absorption;>>Bile secretion;>>Mineral absorption
Gene Name :	ATP1B3
Protein Name :	Sodium/potassium-transporting ATPase subunit beta-3
Human Gene Id :	483
Human Swiss Prot No :	P54709
Mouse Gene Id :	11933
Mouse Swiss Prot No :	P97370
Rat Gene Id :	25390
Rat Swiss Prot No :	Q63377
Immunogen :	The antiserum was produced against synthesized peptide derived from the C-terminal region of human ATP1B3. AA range:222-271
Specificity :	CD298 Polyclonal Antibody detects endogenous levels of CD298 protein.

Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	36kD
Cell Pathway :	Cardiac muscle contraction;Aldosterone-regulated sodium reabsorption;
Background :	<p>The protein encoded by this gene belongs to the family of Na⁺/K⁺ and H⁺/K⁺ ATPases beta chain proteins, and to the subfamily of Na⁺/K⁺ -ATPases. Na⁺/K⁺ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane. The glycoprotein subunit of Na⁺/K⁺ -ATPase is encoded by multiple genes. This gene encodes a beta 3 subunit. This gene encodes a beta 3 subun</p>
Function :	<p>function:This is the non-catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of Na(+) and K(+) ions across the plasma membrane. The exact function of the beta-3 subunit is not known.,similarity:Belongs to the X(+)/potassium ATPases subunit beta family.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Composed of three subunits: alpha (catalytic), beta and gamma.,</p>
Subcellular Location :	Apical cell membrane ; Single-pass type II membrane protein . Basolateral cell membrane ; Single-pass type II membrane protein . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Expression :	Lung,Placenta,Uterus,

Products Images



Western Blot analysis of PC12, NIH-3T3 cells using CD298 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000