



**CRBP II Polyclonal Antibody**

<b>Catalog No :</b>	YT5387
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	CRBP II
<b>Fields :</b>	>>Vitamin digestion and absorption
<b>Gene Name :</b>	RBP2
<b>Protein Name :</b>	Retinol-binding protein 2
<b>Human Gene Id :</b>	5948
<b>Human Swiss Prot No :</b>	P50120
<b>Mouse Gene Id :</b>	19660
<b>Mouse Swiss Prot No :</b>	Q08652
<b>Rat Gene Id :</b>	24710
<b>Rat Swiss Prot No :</b>	P06768
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human RBP2. AA range:71-120
<b>Specificity :</b>	CRBP II Polyclonal Antibody detects endogenous levels of CRBP II protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	15kD
<b>Background :</b>	retinol binding protein 2(RBP2) Homo sapiens This gene encodes an abundant protein present in the small intestinal epithelium. It is thought to participate in the uptake and/or intracellular metabolism of vitamin A. Vitamin A is a fat-soluble vitamin necessary for growth, reproduction, differentiation of epithelial tissues, and vision. This protein may also modulate the supply of retinoic acid to the nuclei of endometrial cells during the menstrual cycle. [provided by RefSeq, Aug 2015],
<b>Function :</b>	domain:Forms a beta-barrel structure that accommodates hydrophobic ligands in its interior.,function:Intracellular transport of retinol.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,tissue specificity:Higher expression in adult small intestine and to a much lesser extent in fetal kidney.,
<b>Subcellular Location :</b>	Cytoplasm.
<b>Expression :</b>	Higher expression in adult small intestine and to a much lesser extent in fetal kidney.

## Products Images





Western blot analysis of lysate from K562 cells, using RBP2 Antibody.