

DNAM-1 Polyclonal Antibody

Catalog No: YT5402

Reactivity: Human; Mouse

Applications: WB;ELISA

Target: DNAM-1

Fields: >>Cell adhesion molecules

Q15762

Q8K4F0

Gene Name: CD226

Protein Name: CD226 antigen

Human Gene Id: 10666

Human Swiss Prot

Tullian Swiss Froi

No:

Mouse Gene ld: 225825

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from the

Internal region of human CD226. AA range:71-120

Specificity: DNAM-1 Polyclonal Antibody detects endogenous levels of DNAM-1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 38kD

Cell Pathway: Cell adhesion molecules (CAMs);

Background: This gene encodes a glycoprotein expressed on the surface of NK cells,

platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Jan 2015],

Function: function: Receptor involved in intercellular adhesion, lymphocyte signaling,

cytotoxicity and lymphokine secretion mediated by cytotoxic T-lymphocyte (CTL)

and NK cell.,PTM:Phosphorylated.,similarity:Contains 2 Ig-like C2-type

(immunoglobulin-like) domains., subunit:Interacts with PVR and PVRL2., tissue


specificity: Expressed by peripheral blood T-lymphocytes.,

Subcellular Location:

Cell membrane ; Single-pass type I membrane protein .


Expression: Expressed by peripheral blood T-lymphocytes.

Products Images



Western Blot analysis of HeLa cells using DNAM-1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





Western blot analysis of lysate from HeLa cells, using CD226 Antibody.