

## CD206 Polyclonal Antibody

<b>Catalog No :</b>	YT5640
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	CD206
<b>Fields :</b>	>>Phagosome;>>Tuberculosis
<b>Gene Name :</b>	MRC1
<b>Protein Name :</b>	Macrophage mannose receptor 1
<b>Human Gene Id :</b>	4360
<b>Human Swiss Prot No :</b>	P22897
<b>Mouse Gene Id :</b>	17533
<b>Mouse Swiss Prot No :</b>	Q61830
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human MRC1. AA range:341-390
<b>Specificity :</b>	CD206 Polyclonal Antibody detects endogenous levels of CD206 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000. IF 1:100-300 Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 170kD


**Background :** The recognition of complex carbohydrate structures on glycoproteins is an important part of several biological processes, including cell-cell recognition, serum glycoprotein turnover, and neutralization of pathogens. The protein encoded by this gene is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. The protein has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment.[provided by RefSeq, Sep 2015],

**Function :** function:Mediates the endocytosis of glycoproteins by macrophages. Binds both sulfated and non-sulfated polysaccharide chains. Acts as phagocytic receptor for bacteria, fungi and other pathogens.,miscellaneous:CRDs 1-3 have at most very weak affinity for carbohydrate. CRD 4 shows the highest affinity binding and has multispecificity for a variety of monosaccharides. At least 3 CRDs (4, 5, and 7) are required for high affinity binding and endocytosis of multivalent glycoconjugates.,online information:Macrophage mannose receptor,similarity:Contains 1 fibronectin type-II domain.,similarity:Contains 1 ricin B-type lectin domain.,similarity:Contains 8 C-type lectin domains.,


**Subcellular Location :** Endosome membrane ; Single-pass type I membrane protein . Cell membrane ; Single-pass type I membrane protein .

**Expression :** Milk,Placenta,Testis,

## Products Images



Western Blot analysis of NIH-3T3 cells using CD206 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western Blot analysis of 3T3 cells using CD206 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000