

## IDH1 Polyclonal Antibody


<b>Catalog No :</b>	YT5416
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	IDH1
<b>Fields :</b>	>>Citrate cycle (TCA cycle);>>Glutathione metabolism;>>Metabolic pathways;>>Carbon metabolism;>>2-Oxocarboxylic acid metabolism;>>Biosynthesis of amino acids;>>Peroxisome;>>Central carbon metabolism in cancer
<b>Gene Name :</b>	IDH1
<b>Protein Name :</b>	Isocitrate dehydrogenase [NADP] cytoplasmic
<b>Human Gene Id :</b>	3417
<b>Human Swiss Prot No :</b>	O75874
<b>Mouse Gene Id :</b>	15926
<b>Mouse Swiss Prot No :</b>	O88844
<b>Rat Gene Id :</b>	24479
<b>Rat Swiss Prot No :</b>	P41562
<b>Immunogen :</b>	Synthesized peptide derived from the N-terminal region of human IDH1.
<b>Specificity :</b>	IDH1 Polyclonal Antibody detects endogenous levels of IDH1 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200

---


<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	46kD
<b>Cell Pathway :</b>	Citrate cycle (TCA cycle);Glutathione metabolism;
<b>Background :</b>	<p>Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to</p>
<b>Function :</b>	<p>catalytic activity:Isocitrate + NADP(+) = 2-oxoglutarate + CO(2) + NADPH.,catalytic activity:Oxalosuccinate + NADP(+) = 2-oxoglutarate + CO(2) + NADPH.,cofactor:Binds 1 magnesium or manganese ion per subunit.,disease:Defects in IDH1 are a cause of glioblastoma multiforme (GBM) [MIM:137800]; also called familial glioma of brain. Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas, glioblastoma multiforme, oligodendrogliomas, and ependymomas.,miscellaneous:Cancer mutations affecting Arg-132 are tissue-specific, and suggest that this residue plays a unique role in the development of high-grade gliomas.,online information:Isocitrate dehydrogenase entry,similarity:Belongs to the isocitrate and isopropylmalate dehydrogenases family.,subunit:Homodimer.,</p>
<b>Subcellular Location :</b>	Cytoplasm, cytosol . Peroxisome .
<b>Expression :</b>	Brain,Cajal-Retzius cell,Fetal brain cortex,Human endometri

---


## Products Images



Western Blot analysis of 293, HeLa cells using IDH1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:100