

LIR-7 Polyclonal Antibody

Catalog No :	YT5466
Reactivity :	Human
Applications :	WB;IHC;IF;ELISA
Target :	LIR-7
Fields :	>>Osteoclast differentiation;>>B cell receptor signaling pathway
Gene Name :	LILRA2
Protein Name :	Leukocyte immunoglobulin-like receptor subfamily A member 2
Human Gene Id :	11027
Human Swiss Prot No :	Q8N149
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human LILRA2. AA range:121-170
Specificity :	LIR-7 Polyclonal Antibody detects endogenous levels of LIR-7 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	53kD

Background :

This gene encodes a member of a family of immunoreceptors that are expressed predominantly on monocytes and B cells, and at lower levels on dendritic cells and natural killer cells. The encoded protein is an activating receptor that inhibits dendritic cell differentiation and antigen presentation and suppresses innate immune response. Alternatively spliced transcript variants encoding different isoforms have been found. This gene is located in a cluster of related genes on chromosome 19 and there is a pseudogene for this gene on chromosome 3. [provided by RefSeq, Mar 2014],

Function :

function:May act as receptor for class I MHC antigens.,similarity:Contains 4 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Expression levels are very low or not detectable on monocytes, T-cells, B-cells, dendritic cells and natural killer (NK) cells.,


Subcellular Location :

Cell membrane ; Single-pass type I membrane protein .; [Isoform 4]: Secreted .


Expression :

Detected on the surface of all peripheral blood monocytes, neutrophils, basophils and eosinophils (at protein level) (PubMed:12529506, PubMed:22479404). Expression levels are very low or not detectable on monocytes, T-cells, B-cells, dendritic cells and natural killer (NK) cells (PubMed:9548455).

Products Images



Western Blot analysis of HepG2 cells using LIR-7 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100