

Stra8 Polyclonal Antibody

Catalog No :	YT5557
Reactivity :	Human;Mouse
Applications :	WB;IHC
Target :	Stra8
Gene Name :	STRA8
Protein Name :	Stimulated by retinoic acid gene 8 protein homolog
Human Gene Id :	346673
Human Swiss Prot No :	Q7Z7C7
Mouse Swiss Prot No :	P70278
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human STRA8. AA range:151-200
Specificity :	Stra8 Polyclonal Antibody detects endogenous levels of Stra8 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	66kD


Background : This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in humans. [provided by RefSeq, Nov 2010],

Function : function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis.,


Subcellular Location : Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent. .

Expression : Expressed specifically in testis and fetal ovaries.

Products Images



Western Blot analysis of NIH-3T3, 293 cells using Stra8 Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).