

Ob Polyclonal Antibody

Catalog No: YT6016

Reactivity: Human; Mouse; Rat

Applications: IHC;IF;ELISA

Target: Ob

Fields: >>Cytokine-cytokine receptor interaction;>>Neuroactive ligand-receptor

interaction;>>AMPK signaling pathway;>>JAK-STAT signaling

pathway;>>Adipocytokine signaling pathway;>>Non-alcoholic fatty liver disease

Gene Name: LEP OB OBS

Protein Name: Leptin (Obese protein) (Obesity factor)

P41159

P41160

Human Gene Id: 3952

Human Swiss Prot

No:

Mouse Gene Id: 16846

Mouse Swiss Prot

No:

Immunogen: Synthetic peptide from human protein at AA range: 10-50

Specificity: The antibody detects endogenous Ob

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : IHC 1:50-200, ELISA 1:10000-20000. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway: Cytokine-cytokine receptor interaction; Neuroactive ligand-receptor

interaction; Jak_STAT; Adipocytokine;

Background: This gene encodes a protein that is secreted by white adipocytes, and which

plays a major role in the regulation of body weight. This protein, which acts through the leptin receptor, functions as part of a signaling pathway that can inhibit food intake and/or regulate energy expenditure to maintain constancy of the adipose mass. This protein also has several endocrine functions, and is involved in the regulation of immune and inflammatory responses, hematopoiesis, angiogenesis and wound healing. Mutations in this gene and/or its regulatory regions cause severe obesity, and morbid obesity with hypogonadism. This gene has also been linked to type 2 diabetes mellitus development. [provided by

RefSeq, Jul 2008],

Function : disease:Defects in LEP may be a cause of autosomal recessive obesity

[MIM:601665].,function:May function as part of a signaling pathway that acts to regulate the size of the body fat depot. An increase in the level of LEP may act directly or indirectly on the CNS to inhibit food intake and/or regulate energy expenditure as part of a homeostatic mechanism to maintain constancy of the adipose mass.,online information:Leptin entry,similarity:Belongs to the leptin

family., subunit: Interacts with SIGLEC6.,

Subcellular Secreted.

Location:

Expression : Adipose tissue is the main source of leptin. It is also produced by other

peripheral tissues such as the skeletal muscle (PubMed:7789654,


PubMed:16052473, PubMed:12448771). Expressed by intercalated and striated

tracts of submandibular and parotid salivary gland intralobular ducts (PubMed:12448771). Detected by fundic epithelium of the gastric mucosa

(PubMed:10896907). Secreted into blood and gastric juice (PubMed:10896907).

Products Images

2/3



Immunohistochemical analysis of paraffin-embedded Human colon. 1, Antibody was diluted at 1:100(4° overnight). 2, Highpressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).