

## TIF-IA Polyclonal Antibody

<b>Catalog No :</b>	YT6132
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC
<b>Target :</b>	TIF-IA
<b>Gene Name :</b>	RRN3 TIFIA
<b>Protein Name :</b>	RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA)
<b>Human Gene Id :</b>	54700
<b>Human Swiss Prot No :</b>	Q9NYV6
<b>Mouse Gene Id :</b>	106298
<b>Mouse Swiss Prot No :</b>	B2RS91
<b>Immunogen :</b>	Synthesized peptide derived from human TIF-IA Polyclonal
<b>Specificity :</b>	This antibody detects endogenous levels of TIF-IA.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 70kD

**Background :** function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,


**Function :** function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,

**Subcellular Location :** Nucleus, nucleolus .


**Location :**

**Expression :** Hippocampus,Kidney,Pooled,Spleen,

## Products Images



Western blot analysis of 3T3 lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).