

IRK14 Polyclonal Antibody

Catalog No :	YN0491
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	IRK14
Fields :	>>Cholinergic synapse;>>Oxytocin signaling pathway
Gene Name :	KCNJ14 IRK4
Protein Name :	ATP-sensitive inward rectifier potassium channel 14 (Inward rectifier K(+) channel Kir2.4) (IRK-4) (Potassium channel, inwardly rectifying subfamily J member 14)
Human Gene Id :	3770
Human Swiss Prot No :	Q9UNX9
Mouse Swiss Prot No :	Q8JZN3
Rat Swiss Prot No :	O70596
Immunogen :	Synthesized peptide derived from human protein . at AA range: 350-430
Specificity :	IRK14 Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Observed Band : 47kD


Background : Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel, and probably has a role in controlling the excitability of motor neurons. [provided by RefSeq, Feb 2013],

Function : function: Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. KCNJ14 gives rise to low-conductance channels with a low affinity to the channel blockers Barium and Cesium., similarity: Belongs to the inward rectifier-type potassium channel family., tissue specificity: Expressed preferentially in retina.,

Subcellular Location : Membrane; Multi-pass membrane protein.

Expression : Expressed preferentially in retina.

Products Images



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°C over night