

## GM130 Polyclonal Antibody

<b>Catalog No :</b>	YT6291
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	GM130
<b>Gene Name :</b>	GOLGA2
<b>Protein Name :</b>	GM130
<b>Human Gene Id :</b>	2801
<b>Human Swiss Prot No :</b>	Q08379
<b>Immunogen :</b>	Synthesized peptide derived from human GM130
<b>Specificity :</b>	This antibody detects endogenous levels of human GM130
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:10000.. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	112-130kD
<b>Background :</b>	The Golgi apparatus, which participates in glycosylation and transport of proteins and lipids in the secretory pathway, consists of a series of stacked cisternae (flattened membrane sacs). Interactions between the Golgi and

microtubules are thought to be important for the reorganization of the Golgi after it fragments during mitosis. This gene encodes one of the golgins, a family of proteins localized to the Golgi. This encoded protein has been postulated to play roles in the stacking of Golgi cisternae and in vesicular transport. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of these variants has not been determined. [provided by RefSeq, Feb 2010],

---

**Function :**

domain:Extended rod-like protein with coiled-coil domains.,function:Golgi auto-antigen; probably involved in maintaining cis-Golgi structure.,sequence caution:Sequence differs from that shown after position 814 due to an internal deletion.,similarity:Belongs to the GOLGA2 family.,subunit:Part of a larger oligomeric complex. Interacts with p115 (By similarity). Interacts with RAB1B that has been activated by GTP-binding. Interacts with GORASP1/GRASP65 and ZFPL1.,

---

**Subcellular Location :**

Golgi apparatus, cis-Golgi network membrane ; Peripheral membrane protein ; Cytoplasmic side . Endoplasmic reticulum-Golgi intermediate compartment membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm, cytoskeleton, spindle pole . Associates with the mitotic spindle during mitosis (PubMed:26165940). .

---

**Expression :**

Brain,Cervix,Colon  
adenocarcinoma,Epithelium,Heart,Hepatocyte,Liver,Placenta,Thymus,

---

## Products Images