

ABCB5 rabbit pAb

Catalog No :	YT7794
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	ABCB5
Fields :	>>ABC transporters
Gene Name :	ABCB5
Protein Name :	ABCB5
Human Gene Id :	340273
Human Swiss Prot No :	Q2M3G0
Immunogen :	Synthesized peptide derived from human ABCB5 AA range: 40-120
Specificity :	This antibody detects endogenous levels of Human ABCB5
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	89kD

Background : ABCB5 belongs to the ATP-binding cassette (ABC) transporter superfamily of integral membrane proteins. These proteins participate in ATP-dependent transmembrane transport of structurally diverse molecules ranging from small ions, sugars, and peptides to more complex organic molecules (Chen et al., 2005 [PubMed 15760339]).[supplied by OMIM, Mar 2008],

Function : caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,caution:Was named ABCB1 by some authors.,function:Plasma membrane transporter able to mediate efflux from cells of the rhodamine dye and of the therapeutic drug doxorubicin. Responsible for the resistance to doxorubicin of a subset of malignant melanomas.,miscellaneous:Depletion of ABCB5 by RNAi increases the sensitivity to several drugs of a subset of melanoma cells.,similarity:Belongs to the ABC transporter family.,similarity:Belongs to the ABC transporter family. Multidrug resistance exporter (TC 3.A.1.201) subfamily.,similarity:Contains 1 ABC transmembrane type-1 domain.,similarity:Contains 1 RRM (RNA recognition motif) domain.,similarity:Contains 2 ABC transporter domains.,subunit:Component of the U11/U12 snRNPs that are part of the U12-type spliceosom

Subcellular Location : Cell membrane ; Multi-pass membrane protein .

Expression : Expressed by CD133-expressing progenitor cells among epidermal melanocytes (at protein level). Widely expressed with specific expression in pigment cells. Highly expressed in several malignant tissues: highly expressed in clinical melanomas, with low expression in normal skin. In melanoma, marks malignant melanoma-initiating cells (MMIC), in which clinical virulence resides as a consequence of unlimited self-renewal capacity, resulting in inexorable tumor progression and metastasis. Also highly expressed in a number of leukemia cells. Expressed in basal limbal epithelium.

Products Images