


RBP rabbit pAb

Catalog No :	YT7861
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC
Target :	RBP
Gene Name :	RBP4 PRO2222
Protein Name :	RBP
Human Gene Id :	5950
Human Swiss Prot No :	P02753
Mouse Gene Id :	19662
Mouse Swiss Prot No :	Q00724
Rat Gene Id :	25703
Rat Swiss Prot No :	P04916
Immunogen :	Synthesized peptide derived from human RBP AA range: 11-60
Specificity :	This antibody detects endogenous levels of Human RBP
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	<u>1 mg/ml</u>
Storage Stability :	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
Molecularweight :	<u>22kD</u>
Background :	<u>retinol binding protein 4(RBP4) Homo sapiens This protein belongs to the lipocalin family and is the specific carrier for retinol (vitamin A alcohol) in the blood. It delivers retinol from the liver stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin which prevents its loss by filtration through the kidney glomeruli. A deficiency of vitamin A blocks secretion of the binding protein posttranslationally and results in defective delivery and supply to the epidermal cells. [provided by RefSeq, Jul 2008],</u>
Function :	<u>disease:A deficiency of vitamin A blocks secretion of the binding protein post-translationally and results in defective delivery and supply of vitamin to the epidermal cells (a condition associated with a dermatosis).,disease:Defects in RBP4 are a cause of retinol-binding protein deficiency [MIM:180250]. This condition causes night vision problems. It produces a typical "fundus xerophthalmicus," featuring a progressed atrophy of the retinal pigment epithelium.,function:Delivers retinol from the liver stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin, this prevents its loss by filtration through the kidney glomeruli.,mass spectrometry: PubMed:12237133,mass spectrometry: PubMed:7666002,online information:Retina International's Scientific Newsletter,online information:Retinol-binding protein 4 entry,similarity:Belongs to the calycin superfamil</u>
Subcellular Location :	<u>Secreted .</u>
Expression :	<u>Detected in blood plasma and in urine (at protein level).</u>

Products Images



Immunohistochemical analysis of paraffin-embedded human Breast cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).