

WNT7A Polyclonal Antibody

Catalog No :	YN0287
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	WNT7A
Fields :	>>mTOR signaling pathway;>>Wnt signaling pathway;>>Hippo signaling pathway;>>Signaling pathways regulating pluripotency of stem cells;>>Melanogenesis;>>Cushing syndrome;>>Alzheimer disease;>>Pathways of neurodegeneration - multiple diseases;>>Human papillomavirus infection;>>Pathways in cancer;>>Proteoglycans in cancer;>>Basal cell carcinoma;>>Breast cancer;>>Hepatocellular carcinoma;>>Gastric cancer
Gene Name :	WNT7A
Protein Name :	Protein Wnt-7a
Human Gene Id :	7476
Human Swiss Prot No :	O00755
Mouse Swiss Prot No :	P24383
Immunogen :	Synthesized peptide derived from human protein . at AA range: 110-190
Specificity :	WNT7A Polyclonal Antibody detects endogenous levels of protein.
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 ELISA 1:5000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 38kD

Cell Pathway : WNT;WNT-T CELLHedgehog;Melanogenesis;Pathways in cancer;Basal cell carcinoma;


Background : This gene is a member of the WNT gene family, which consists of structurally related genes that encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is involved in the development of the anterior-posterior axis in the female reproductive tract, and also plays a critical role in uterine smooth muscle patterning and maintenance of adult uterine function. Mutations in this gene are associated with Fuhrmann and Al-Awadi/Raas-Rothschild/Schinzel phocomelia syndromes. [provided by RefSeq, Jul 2008],

Function : disease:Defects in WNT7A are a cause of Fuhrmann syndrome [MIM:228930]; also called fibular aplasia or hypoplasia femoral bowing and poly- syn- and oligodactyly. Fuhrmann syndrome is a distinct limb-malformation disorder characterized also by various degrees of limb aplasia/hypoplasia and joint dysplasia.,disease:Defects in WNT7A are the cause of limb/pelvis-hypoplasia/aplasia syndrome (LPHAS) [MIM:276820]; also called absence of ulna and fibula with severe limb deficiency. LPHAS is a limb-malformation disorder characterized by various degrees of limb aplasia/hypoplasia and joint dysplasia.,function:Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. Signaling by Wnt-7a allows sexually dimorphic development of the mullerian ducts.,similarity:Belongs to the Wnt family.,subunit:Interacts with PORCN.,tissue specificity:Expression is re

Subcellular Location : Secreted, extracellular space, extracellular matrix . Secreted .

Expression : Expression is restricted to placenta, kidney, testis, uterus, fetal lung, and fetal and adult brain.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night