


Dynamin I (phospho Ser774) Polyclonal Antibody


Catalog No :	YP0627
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Dynamin I
Fields :	>>Phospholipase D signaling pathway;>>Endocytosis;>>Synaptic vesicle cycle;>>Endocrine and other factor-regulated calcium reabsorption;>>Bacterial invasion of epithelial cells
Gene Name :	DNM1
Protein Name :	Dynamin-1
Human Gene Id :	1759
Human Swiss Prot No :	Q05193
Mouse Gene Id :	13429
Mouse Swiss Prot No :	P39053
Rat Gene Id :	140694
Rat Swiss Prot No :	P21575
Immunogen :	The antiserum was produced against synthesized peptide derived from human Dynamin-1 around the phosphorylation site of Ser774. AA range:740-789
Specificity :	Phospho-Dynamin I (S774) Polyclonal Antibody detects endogenous levels of Dynamin I protein only when phosphorylated at S774.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	97kD
Cell Pathway :	Endocytosis;Fc gamma R-mediated phagocytosis;
Background :	<p>dynamamin 1(DNM1) Homo sapiens This gene encodes a member of the dynamamin subfamily of GTP-binding proteins. The encoded protein possesses unique mechanochemical properties used to tubulate and sever membranes, and is involved in clathrin-mediated endocytosis and other vesicular trafficking processes. Actin and other cytoskeletal proteins act as binding partners for the encoded protein, which can also self-assemble leading to stimulation of GTPase activity. More than sixty highly conserved copies of the 3' region of this gene are found elsewhere in the genome, particularly on chromosomes Y and 15. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],</p>
Function :	<p>catalytic activity:GTP + H₂O = GDP + phosphate.,function:Microtubule-associated force-producing protein involved in producing microtubule bundles and able to bind and hydrolyze GTP. Most probably involved in vesicular trafficking processes, in particular endocytosis.,similarity:Belongs to the dynamamin family.,similarity:Contains 1 GED domain.,similarity:Contains 1 PH domain.,subcellular location:Microtubule-associated.,subunit:Interacts with CAV1 and SH3GLB1. Binds SH3GL1, SH3GL2 and SH3GL3.,</p>
Subcellular Location :	Cytoplasm . Cytoplasm, cytoskeleton . Microtubule-associated.
Expression :	Brain,Platelet,PNS,


Products Images



Western Blot analysis of mouse cells using Phospho-Dynamin I (S774) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using Dynamin-1 (Phospho-Ser774) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from mouse brain, using Dynamin-1 (Phospho-Ser774) Antibody. The lane on the right is blocked with the phospho peptide.