

## Fyn (phospho Tyr530) Polyclonal Antibody

Catalog No: YP0632

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;IHC;IF;ELISA

Target: Fyn

**Fields:** >>Sphingolipid signaling pathway;>>Phospholipase D signaling

pathway;>>Axon guidance;>>Osteoclast differentiation;>>Focal

adhesion;>>Adherens junction;>>Platelet activation;>>Natural killer cell mediated

cytotoxicity;>>T cell receptor signaling pathway;>>Fc epsilon RI signaling pathway;>>Cholinergic synapse;>>Prion disease;>>Pathogenic Escherichia coli

infection;>>Viral myocarditis

Gene Name: FYN

**Protein Name:** Tyrosine-protein kinase Fyn

P06241

P39688

Human Gene Id: 2534

**Human Swiss Prot** 

No:

Mouse Gene Id: 14360

**Mouse Swiss Prot** 

No:

Rat Gene ld: 25150

Rat Swiss Prot No: Q62844

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

Fyn around the phosphorylation site of Tyr530. AA range:488-537

**Specificity:** Phospho-Fyn (Y530) Polyclonal Antibody detects endogenous levels of Fyn

protein only when phosphorylated at Y530.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

1/3



**Source :** Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 60kD

Cell Pathway: Axon guidance; Focal adhesion; Adherens\_Junction; Natural killer cell mediated

cytotoxicity; T Cell Receptor; Fc epsilon RI; Prion diseases; Pathogenic

Escherichia coli infection; Viral myocarditis;

**Background:** This gene is a member of the protein-tyrosine kinase oncogene family. It

encodes a membrane-associated tyrosine kinase that has been implicated in the

control of cell growth. The protein associates with the p85 subunit of phosphatidylinositol 3-kinase and interacts with the fvn-binding protein.

Alternatively spliced transcript variants encoding distinct isoforms exist. [provided

by RefSeq, Jul 2008],

**Function :** catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine

phosphate.,cofactor:Manganese.,enzyme regulation:Inhibited by phosphorylation of Tyr-531 by leukocyte common antigen and activated by dephosphorylation of

this site., function: Implicated in the control of cell growth. Plays a role in the

regulation of intracellular calcium levels, with isoform 2 showing the greater ability to mobilize cytoplasmic calcium in comparison to isoform 1. Required in brain development and mature brain function with important roles in the regulation of axon growth, axon guidance, and neurite extension. Blocks axon outgrowth and

attraction induced by NTN1 by phosphorylating its receptor

DDC.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase

family. SRC subfamily., similarity: Contains 1


Subcellular Cytoplasm. Nucleus. Cell membrane. Present and active in lipid rafts.

**Location :** Palmitoylation is crucial for proper trafficking.


**Expression:** Isoform 1 is highly expressed in the brain. Isoform 2 is expressed in cells of

hemopoietic lineages, especially T-lymphocytes.


## **Products Images**



Western Blot analysis of LOVO+UV cells using Phospho-Fyn (Y530) Polyclonal Antibody diluted at 1:2000



Western blot analysis of 3T3 lysis using Phospho-Fyn (Y530) antibody. Antibody was diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Fyn (Phospho-Tyr530) Antibody. The picture on the right is blocked with the phospho peptide.