


PAK β (phospho Ser154) Polyclonal Antibody


Catalog No :	YP0795
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PAK β
Fields :	>>ErbB signaling pathway;>>Ras signaling pathway;>>Axon guidance;>>Focal adhesion;>>T cell receptor signaling pathway;>>Regulation of actin cytoskeleton;>>Pathogenic Escherichia coli infection;>>Salmonella infection;>>Human immunodeficiency virus 1 infection;>>Renal cell carcinoma
Gene Name :	PAK3
Protein Name :	Serine/threonine-protein kinase PAK 3
Human Gene Id :	5063
Human Swiss Prot No :	O75914
Mouse Gene Id :	18481
Mouse Swiss Prot No :	Q61036
Rat Gene Id :	29433
Rat Swiss Prot No :	Q62829
Immunogen :	The antiserum was produced against synthesized peptide derived from human PAK3 around the phosphorylation site of Ser154. AA range:121-170
Specificity :	Phospho-PAK β (S154) Polyclonal Antibody detects endogenous levels of PAK β protein only when phosphorylated at S154.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	<u>WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage Stability :	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
Observed Band :	<u>72kD</u>
Cell Pathway :	<u>ErbB_HER;Axon guidance;Focal adhesion;T_Cell_Receptor;Regulates Actin and Cytoskeleton;Renal cell carcinoma;</u>
Background :	<u>The protein encoded by this gene is a serine-threonine kinase and forms an activated complex with GTP-bound RAS-like (P21), CDC2 and RAC1. This protein may be necessary for dendritic development and for the rapid cytoskeletal reorganization in dendritic spines associated with synaptic plasticity. Defects in this gene are the cause of non-syndromic mental retardation X-linked type 30 (MRX30), also called X-linked mental retardation type 47 (MRX47). Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Apr 2016],</u>
Function :	<u>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,disease:Defects in PAK3 are the cause of mental retardation X-linked type 30 (MRX30) [MIM:300558]; also called X-linked mental retardation type 47 (MRX47). Mental retardation is a mental disorder characterized by significantly sub-average general intellectual functioning associated with impairments in adaptative behavior and manifested during the developmental period. Non-syndromic mental retardation patients do not manifest other clinical signs.,enzyme regulation:Activated by binding small G proteins. Binding of GTP-bound CDC42 or RAC1 to the autoregulatory region releases monomers from the autoinhibited dimer, enables phosphorylation of Thr-436 and allows the kinase domain to adopt an active structure.,function:Key regulator of synapse formation and plasticity in the hippocampus.,PTM:Autophosphorylated whe</u>
Subcellular Location :	<u>Cytoplasm .</u>
Expression :	<u>Restricted to the nervous system. Highly expressed in postmitotic neurons of the developing and postnatal cerebral cortex and hippocampus.</u>


Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PAK3 (Phospho-Ser154) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using PAK3 (Phospho-Ser154) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from rat heart, using PAK3 (Phospho-Ser154) Antibody. The lane on the right is blocked with the phospho peptide.