


## Rho GDIa (phospho Ser174) Polyclonal Antibody

Catalog No :	YP0762
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Rho GDIa
Fields :	>>Neurotrophin signaling pathway;>>Vasopressin-regulated water reabsorption
Gene Name :	ARHGDIA
Protein Name :	Rho GDP-dissociation inhibitor 1
Human Gene Id :	396
Human Swiss Prot	P52565
No : Mouse Gene Id :	192662
Mouse Swiss Prot	Q99PT1
No : Rat Gene Id :	360678
Rat Swiss Prot No :	Q5XI73
Immunogen :	The antiserum was produced against synthesized peptide derived from human ARHGDIA around the phosphorylation site of Ser174. AA range:140-189
Specificity :	Phospho-Rho GDIa (S174) Polyclonal Antibody detects endogenous levels of Rho GDIa protein only when phosphorylated at S174.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000 IF 1:50-200



	,
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-
	chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
oonoonnation .	
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	23kD
Cell Pathway :	Neurotrophin;
Deelverreund :	This gaps appedes a protein that plays a key role in the regulation of signaling
Background :	This gene encodes a protein that plays a key role in the regulation of signaling through Rho GTPases. The encoded protein inhibits the disassociation of Rho
	family members from GDP (guanine diphosphate), thereby maintaining these
	factors in an inactive state. Activity of this protein is important in a variety of
	cellular processes, and expression of this gene may be altered in tumors.
	Mutations in this gene have been found in individuals with nephrotic syndrome,
	type 8. Alternate splicing results in multiple transcript variants. [provided by
	RefSeq, Jul 2014],
Function :	function:Regulates the GDP/GTP exchange reaction of the Rho proteins by
	inhibiting the dissociation of GDP from them, and the subsequent binding of GTP
	to them.,similarity:Belongs to the Rho GDI family.,subunit:Monomer.,
Subcellular	Cytoplasm .
Location :	
Expression :	Brain,Cajal-Retzius cell,Colon,Epithelium,Fetal brain cortex,Lung,M




## **Products Images**

Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using ARHGDIA (Phospho-Ser174) Antibody



Immunohistochemistry analysis of paraffin-embedded human cervix carcinoma, using ARHGDIA (Phospho-Ser174) Antibody. The picture on the right is blocked with the phospho peptide.



-- 19 (kD) Western blot analysis of lysates from HUVEC cells treated with EGF 200ng/ml 30', using ARHGDIA (Phospho-Ser174) Antibody. The lane on the right is blocked with the phospho peptide.