

p90RSK (Phospho Thr359/Ser363) rabbit pAb

Catalog No: YP1428

Reactivity: Human; Mouse; Rat

Applications: WB

Target: Rsk-1

Fields: >>MAPK signaling pathway;>>Oocyte meiosis;>>mTOR signaling

pathway;>>Thermogenesis;>>Long-term potentiation;>>Neurotrophin signaling

pathway;>>Progesterone-mediated oocyte maturation;>>Insulin

resistance;>>Yersinia infection;>>Chemical carcinogenesis - receptor activation

Gene Name: RPS6KA1 MAPKAPK1A RSK1

Q15418

P18653

Protein Name: p90RSK (Thr359/Ser363)

Human Gene Id: 6195

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Rat Gene Id: 81771

Rat Swiss Prot No: Q63531

Immunogen: Synthesized phosho peptide around human p90RSK (Thr359 and Ser363)

Specificity: This antibody detects endogenous levels of Human Mouse Rat p90RSK

(phospho-Thr359 or Ser363)

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:1000-2000

1/2



Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 83kD

Cell Pathway: Regulates Angiogenesis; Insulin Receptor; B Cell Receptor; AMPK

Background: ribosomal protein S6 kinase A1(RPS6KA1) Homo sapiens This gene encodes a

member of the RSK (ribosomal S6 kinase) family of serine/threonine kinases. This kinase contains 2 nonidentical kinase catalytic domains and phosphorylates various substrates, including members of the mitogen-activated kinase (MAPK) signalling pathway. The activity of this protein has been implicated in controlling cell growth and differentiation. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],

Function: catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:The

sequence shown here is derived from an Ensembl automatic analysis pipeline and

should be considered as preliminary data.,cofactor:Magnesium.,enzyme regulation:Activated by multiple phosphorylations on threonine and serine residues.,function:Serine/threonine kinase that may play a role in mediating the

growth-factor and stress induced activation of the transcription factor CREB.,PTM:Autophosphorylated on Ser-380, as part of the activation

process.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase

subfamily., similarity: Contains 1 AGC-kinase C-terminal

domain., similarity: Contains 2 protein kinase domains., subunit: Forms a complex with either ERK1 or ERK2 in quiescent cells. Transiently dissociates following

mitogenic s

Subcellular Location:

Nucleus. Cytoplasm.

Expression:

Colon, Epithelium,

Products Images