

## TAZ (Phospho Ser89) rabbit pAb

YP1526 Catalog No:

Human; Mouse; Rat Reactivity:

**Applications:** WB

Target: TAZ

Fields: >>Hippo signaling pathway;>>Hippo signaling pathway - multiple species

**Gene Name:** WWTR1 TAZ

**Protein Name:** TAZ (Ser89)

**Human Gene Id:** 25937

**Human Swiss Prot** 

Q9GZV5

No:

Mouse Gene Id: 97064

**Mouse Swiss Prot** 

No:

Synthesized phosho peptide around human TAZ (Ser89) Immunogen:

This antibody detects endogenous levels of Human Mouse Rat TAZ (phospho-**Specificity:** 

Ser89)

Q9EPK5

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

WB 1:1000-2000 **Dilution:** 

**Purification:** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

**Concentration:** 1 mg/ml

1/2

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

44kD

Observed Band:

#### **Function:**

domain:Binds to transcription factors via its WW domain.,domain:The PDZ-binding motif is essential for stimulated gene transcription. It localizes the protein into both punctate nuclear foci and plasma membrane-associated complexes.,function:Functions as a transcriptional coactivator.,PTM:Phosphorylated. Phosphorylation results in the inhibition of transcriptional coactivation through YWHAZ-mediated nuclear export.,similarity:Contains 1 WW domain.,subcellular location:Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies. When phosphorylated, is retained in cytoplasm by YWHAZ.,subunit:Binds to SLC9A3R2 via the PDZ motif at the plasma membrane. Binds to YWHAZ in vivo and in vitro through the phosphoserine-binding motif RSHSSP.,tissue specificity:Highly expressed in kidney, heart, placenta and lung.,


# Subcellular Location:

Nucleus . Cytoplasm . Cell membrane . Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:18568018). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:21145499). At blastocyst stage expressed in the nucleus in trophectodermal cells, however expressed in the cytoplasm in the inner cell mass (By similarity). .

### **Expression:**

Highly expressed in kidney, heart, placenta and lung. Expressed in the thyroid tissue.

## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night