

**p95/NBS1 (Phospho Ser343) rabbit pAb**

<b>Catalog No :</b>	YP1430
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB
<b>Target :</b>	p95/NBS1
<b>Fields :</b>	>>Homologous recombination;>>Cellular senescence
<b>Gene Name :</b>	NBN NBS NBS1 P95
<b>Protein Name :</b>	p95/NBS1 (Ser343)
<b>Human Gene Id :</b>	4683
<b>Human Swiss Prot No :</b>	O60934
<b>Mouse Gene Id :</b>	27354
<b>Mouse Swiss Prot No :</b>	Q9R207
<b>Rat Gene Id :</b>	85482
<b>Rat Swiss Prot No :</b>	Q9JIL9
<b>Immunogen :</b>	Synthesized phospho peptide around human p95 (Ser343)
<b>Specificity :</b>	This antibody detects endogenous levels of Human Mouse p95 or NBS1 (phospho-Ser343)
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:1000-2000

---

<b>Purification :</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	85kD
<b>Cell Pathway :</b>	Homologous recombination;
<b>Background :</b>	Mutations in this gene are associated with Nijmegen breakage syndrome, an autosomal recessive chromosomal instability syndrome characterized by microcephaly, growth retardation, immunodeficiency, and cancer predisposition. The encoded protein is a member of the MRE11/RAD50 double-strand break repair complex which consists of 5 proteins. This gene product is thought to be involved in DNA double-strand break repair and DNA damage-induced checkpoint activation. [provided by RefSeq, Jul 2008],
<b>Function :</b>	disease:Defects in NBN are a cause of genetic susceptibility to breast cancer (BC) [MIM:114480]. BC is an extremely common malignancy, affecting one in eight women during their lifetime. A positive family history has been identified as major contributor to risk of development of the disease, and this link is striking for early-onset breast cancer.,disease:Defects in NBN are the cause of Nijmegen breakage syndrome (NBS) [MIM:251260]. NBS is an autosomal recessive syndrome characterized by chromosomal instability, radiation sensitivity, microcephaly, growth retardation, immunodeficiency and predisposition to cancer, particularly to lymphoid malignancies.,disease:Defects in NBN may be associated with aplastic anemia [MIM:609135]. Aplastic anemia is a disease of bone-marrow failure characterized by peripheral pancytopenia and marrow hypoplasia. Most of the cases of aplastic anemia are idiopa
<b>Subcellular Location :</b>	Nucleus . Nucleus, PML body . Chromosome, telomere . Chromosome . Localizes to discrete nuclear foci after treatment with genotoxic agents (PubMed:26438602, PubMed:10783165, PubMed:26215093). Acetylation of 'Lys-5' of histone H2AX (H2AXK5ac) promotes NBN/NBS1 assembly at the sites of DNA damage (PubMed:26438602). .
<b>Expression :</b>	Ubiquitous (PubMed:9590180). Expressed at high levels in testis (PubMed:9590180).

---

## Products Images