

EIF6 (Phospho Ser235) rabbit pAb

Catalog No: YP1728

Reactivity: Human; Mouse; Rat

Applications: WB

Target: EIF6

Fields: >>Ribosome biogenesis in eukaryotes

Gene Name: EIF6 EIF3A ITGB4BP OK/SW-cl.27

Protein Name: EIF6 (Phospho-Ser235)

Human Gene Id: 3692

Human Swiss Prot

P56537

No:

Mouse Gene Id: 16418

Mouse Swiss Prot

O55135

No:

Rat Gene Id: 305506

Rat Swiss Prot No: Q3KRD8

Immunogen: Synthesized peptide derived from human EIF6 (Phospho-Ser235)

Specificity: This antibody detects endogenous levels of EIF6 (Phospho-Ser235) at Human,

Mouse,Rat

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500-2000

1/2



Purification: The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 27kD

Background: Hemidesmosomes are structures which link the basal lamina to the intermediate

filament cytoskeleton. An important functional component of hemidesmosomes is the integrin beta-4 subunit (ITGB4), a protein containing two fibronectin type III domains. The protein encoded by this gene binds to the fibronectin type III domains of ITGB4 and may help link ITGB4 to the intermediate filament cytoskeleton. The encoded protein, which is insoluble and found both in the nucleus and in the cytoplasm, can function as a translation initiation factor and prevent the association of the 40S and 60S ribosomal subunits. Multiple non-protein coding transcript variants and variants encoding two different isoforms

have been found for this gene. [provided by RefSeg, Jun 2012].

Function: function:Binds to the 60S ribosomal subunit and prevents its association with the

40S ribosomal subunit to form the 80S initiation complex., similarity: Belongs to the

eIF-6 family.,subunit:Monomer.,

Subcellular Cytoplasm. Nucleus, nucleolus. Shuttles between cytoplasm and

Location: nucleus/nucleolus.

Expression: Expressed at very high levels in colon carcinoma with lower levels in normal

colon and ileum and lowest levels in kidney and muscle (at protein level).

Products Images





Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000