

RUNX1 (Acetyl Lys24) rabbit pAb

Catalog No :	YK0098
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	RUNX1
Fields :	>>Tight junction;>>Th17 cell differentiation;>>Pathways in cancer;>>Transcriptional misregulation in cancer;>>Chronic myeloid leukemia;>>Acute myeloid leukemia
Gene Name :	RUNX1 AML1 CBFA2
Protein Name :	RUNX1 (Acetyl Lys24)
Human Gene Id :	861
Human Swiss Prot No :	Q01196
Mouse Gene Id :	12394
Mouse Swiss Prot No :	Q03347
Rat Gene Id :	50662
Rat Swiss Prot No :	Q63046
Immunogen :	Synthesized peptide derived from human RUNX1 (Acetyl Lys24)
Specificity :	This antibody detects endogenous levels of Human,Mouse,Rat RUNX1 (Acetyl Lys24)
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:1000-2000 ELISA 1:5000-20000

Purification : The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 50kD

Background : alternative products:Additional isoforms seem to exist,caution:The fusion of AML1 with EAP in T-MDS induces a change of reading frame in the latter resulting in 17 AA unrelated to those of EAP.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of chronic myelogenous leukemia (CML). Translocation t(3;21)(q26;q22) with EAP, MSD1 or EVI1.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of chronic myelomonocytic leukemia. Inversion inv(21)(q21;q22) with USP16.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of M2 type acute myeloid leukemia (AML-M2). Translocation t(8;21)(q22;q22) with RUNX1T1/MTG8/ETO.,disease:A chromosomal aberration involving RUNX1/AML1 is a cause of therapy-related myelodysplastic syndrome (T-MDS). Translocation t(3;21)(q26;q22) with EAP, MSD1 or EVI1.,disease:A chromosomal aberration involving RUNX1/AML1 is found in childhood acute lymphoblastic leukemia (ALL). Translocation t(12;21)(p13;q22) with TEL. The translocation fuses the 3'-end of TEL to the alternate 5'-exon of AML-1H.,disease:A chromosomal aberration involving RUNX1/AML1 is found in therapy-related myeloid malignancies. Translocation t(16;21)(q24;q22) that forms a RUNX1-CBFA2T3 fusion protein.,disease:Defects in RUNX1 are the cause of familial platelet disorder with associated myeloid malignancy (FPDMM) [MIM:601399]. FPDMM is an autosomal dominant disease characterized by qualitative and quantitative platelet defects, and propensity to develop acute myelogenous leukemia.,domain:A proline/serine/threonine rich region at the C-terminus is necessary for transcriptional activation of target genes.,function:CBF binds to the core site, 5'-PYGPYGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, LCK, IL-3 and GM-CSF promoters. The alpha subunit binds DNA and appears to have a role in the development of normal hematopoiesis. Isoform AML-1L interferes with the transactivation activity of RUNX1. Acts synergistically with ELF4 to transactivate the IL-3 promoter and with ELF2 to transactivate the mouse BLK promoter. Inhibits MYST4-dependent transcriptional activation.,PTM:Methylated.,PTM:Phosphorylated in its C-terminus upon IL-6 treatment. Phosphorylation enhances interaction with MYST3.,similarity:Contains 1 Runt domain.,subunit:Heterodimer with CBFβ. RUNX1 binds DNA as a monomer and through the Runt domain. DNA-binding is increased by heterodimerization. Isoform AML-1L can neither bind DNA nor heterodimerize. Interacts with TLE1 and THOC4. Interacts with ELF1, ELF2 and SPI1. Interacts via its Runt domain with the ELF4 N-terminal region. Interaction with ELF2 isoform 2 (NERF-1a) may act to repress RUNX1-mediated transactivation. Interacts with MYST3 and MYST4. Interacts with SUV39H1, leading to abrogate the transactivating and DNA-binding properties of RUNX1.,tissue specificity:Expressed in all tissues examined except brain and

heart. Highest levels in thymus, bone marrow and peripheral blood.,

Function :

immune system development, regulation of myeloid leukocyte differentiation, negative regulation of myeloid leukocyte differentiation, positive regulation of myeloid leukocyte differentiation, transcription, regulation of transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter, positive regulation of biosynthetic process, positive regulation of macromolecule biosynthetic process, positive regulation of macromolecule metabolic process, positive regulation of gene expression, hemopoiesis, myeloid cell differentiation, regulation of granulocyte differentiation, negative regulation of granulocyte differentiation, positive regulation of granulocyte differentiation, positive regulation of cellular biosynthetic process, regulation of transcription, negative regulation of cell differentiation, positive regulation of cell differentiation, regulation of myeloid cell d

Subcellular Location :

Nucleus.

Expression :

Expressed in all tissues examined except brain and heart. Highest levels in thymus, bone marrow and peripheral blood.

Products Images