

CD326/EPCAM (PN0555) Nb-FC recombinant antibody

Catalog No :	YA0295
Reactivity :	Human
Applications :	ELISA
Target :	CD326/EPCAM
Gene Name :	EPCAM GA733-2 M1S2 M4S1 MIC18 TACSTD1 TROP1
Protein Name :	Epithelial cell adhesion molecule (Ep-CAM) (Adenocarcinoma-associated antigen) (Cell surface glycoprotein Trop-1) (Epithelial cell surface antigen) (Epithelial glycoprotein) (EGP) (Epithelial glycopro
Human Gene Id :	4072
Human Swiss Prot No :	P16422
Immunogen :	Purified recombinant Human CD326
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of CD326/EPCAM protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse $\mbox{IgG1}$ Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on



	most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations inThis gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008]
Subcellular Location :	Lateral cell membrane ; Single-pass type I membrane protein . Cell junction, tight junction . Colocalizes with CLDN7 at the lateral cell membrane and tight junction
Expression :	Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC). Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Products Images