

CD48 (PN0201) Nb-FC recombinant antibody

Catalog No: YA0386

Reactivity: Human

Applications: ELISA;FCM

Target: CD48

Gene Name: CD48 BCM1 BLAST1

Protein Name : CD48 antigen (B-lymphocyte activation marker BLAST-1) (BCM1 surface

antigen) (Leukocyte antigen MEM-102) (SLAM family member 2) (SLAMF2)

(Signaling lymphocytic activation molecule 2) (TCT.1) (CD antig

Human Gene Id: 962

Human Swiss Prot

No:

Immunogen: Purified recombinant Human CD48

P09326

Specificity: This recombinant monoclonal antibody can detects endogenous levels of CD48

protein.

Formulation : Phosphate-buffered solution

Source: Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain,

recombinantly produced from 293F cell

Dilution: ELISA 1:5000-100000 FCM 1-2µg/Test

Purification: Recombinant Expression and Affinity purified

Concentration: Please check the information on the tube

Storage Stability: -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

Cell Pathway : Natural killer cell mediated cytotoxicity;

1/2



Background:

This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found forThis gene. [provided by RefSeq, Dec 2011]

Function:

Ligand for CD2. Might facilitate interaction between activated lymphocytes. Probably involved in regulating T-cell activation.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,

Subcellular Location:

Cell membrane; Lipid-anchor, GPI-anchor. Secreted.

Expression: Widely express

Widely expressed on all hematopoietic cells.

Products Images

