

**CD69 (PN0005) Nb-FC recombinant antibody**

<b>Catalog No :</b>	YA0466
<b>Reactivity :</b>	Human
<b>Applications :</b>	ELISA
<b>Target :</b>	CD69
<b>Gene Name :</b>	CD69 CLEC2C
<b>Protein Name :</b>	Early activation antigen CD69 (Activation inducer molecule) (AIM) (BL-AC/P26) (C-type lectin domain family 2 member C) (EA1) (Early T-cell activation antigen p60) (GP32/28) (Leukocyte surface antigen)
<b>Human Gene Id :</b>	969
<b>Human Swiss Prot No :</b>	Q07108
<b>Immunogen :</b>	Purified recombinant Human CD69
<b>Specificity :</b>	This recombinant monoclonal antibody can detects endogenous levels of CD69 protein.
<b>Formulation :</b>	Phosphate-buffered solution
<b>Source :</b>	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
<b>Dilution :</b>	ELISA 1:5000-100000
<b>Purification :</b>	Recombinant Expression and Affinity purified
<b>Concentration :</b>	Please check the information on the tube
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
<b>Background :</b>	This gene encodes a member of the calcium dependent lectin superfamily of type II transmembrane receptors. Expression of the encoded protein is induced

upon activation of T lymphocytes, and may play a role in proliferation. Furthermore, the protein may act to transmit signals in natural killer cells and platelets. [provided by RefSeq, Aug 2011]

**Function :**

developmental stage:Earliest inducible cell surface glycoprotein acquired during lymphoid activation.,Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets.,induction:By antigens, mitogens or activators of PKC on the surface of T and B-lymphocytes. By interaction of IL-2 with the p75 IL-2R on the surface of NK cells.,online information:CD69,PTM:Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes.,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer; disulfide-linked.,tissue specificity:Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets.,

**Subcellular Location :**

Membrane; Single-pass type II membrane protein.

**Expression :**

Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets.

## Products Images

