

NKp80 (PN0292) Nb-FC recombinant antibody

Catalog No :	YA0638
Reactivity :	Human
Applications :	ELISA;FCM
Target :	NKp80
Gene Name :	KLRF1 CLEC5C ML
Protein Name :	Killer cell lectin-like receptor subfamily F member 1 (Lectin-like receptor F1) (Activating coreceptor NKp80) (C-type lectin domain family 5 member C)
Human Gene Id :	51348
Human Swiss Prot No :	Q9NZS2
Immunogen :	Purified recombinant Human NKp80
Specificity :	This recombinant monoclonal antibody can detects endogenous levels of NKp80 protein.
Formulation :	Phosphate-buffered solution
Source :	Camel, chimeric fusion of Nanobody (VHH) and mouse IgG1 Fc domain , recombinantly produced from 293F cell
Dilution :	ELISA 1:5000-100000 FCM 1-2µg/Test
Purification :	Recombinant Expression and Affinity purified
Concentration :	Please check the information on the tube
Storage Stability :	-15°C to -25°C/1 year(Avoid freeze / thaw cycles)
Background :	KLRF1, an activating homodimeric C-type lectin-like receptor (CLR), is expressed on nearly all natural killer (NK) cells and stimulates their cytotoxicity and cytokine release (Kuttruff et al., 2009 [PubMed 18922855]).[supplied by OMIM,

Oct 2009]

Function :

Involved in the natural killer (NK)-mediated cytotoxicity of PHA-induced lymphoblasts.,online information:KLRF1,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer.,tissue specificity:Strongly expressed in peripheral blood leukocytes and spleen, with weaker expression in lymph node and adult liver, and no expression detected in bone marrow, thymus, and fetal liver. Not expressed in brain, heart, placenta, lung, kidney, skeletal muscle, and pancreas. Within peripheral blood leukocyte and immunocyte cell lines, expression was predominant in NK cells but was also detected in monocytes.,

Subcellular Location :

Membrane ; Single-pass type II membrane protein .

Expression :

Strongly expressed in peripheral blood leukocytes and spleen, with weaker expression in lymph node and adult liver, and no expression detected in bone marrow, thymus, and fetal liver. Not expressed in brain, heart, placenta, lung, kidney, skeletal muscle, and pancreas. Within peripheral blood leukocyte and immunocyte cell lines, expression was predominant in NK cells but was also detected in monocytes.

Products Images

