

BLNK Monoclonal Antibody

| | |
|------------------------------|---|
| Catalog No : | YM0068 |
| Reactivity : | Human;Mouse |
| Applications : | WB;IHC;IF;FCM;ELISA |
| Target : | BLNK |
| Fields : | >>NF-kappa B signaling pathway;>>Osteoclast differentiation;>>B cell receptor signaling pathway;>>Epstein-Barr virus infection;>>Primary immunodeficiency |
| Gene Name : | BLNK |
| Protein Name : | B-cell linker protein |
| Human Gene Id : | 29760 |
| Human Swiss Prot No : | Q8WV28 |
| Mouse Gene Id : | 17060 |
| Mouse Swiss Prot No : | Q9QUN3 |
| Immunogen : | Purified recombinant fragment of human BLNK expressed in E. Coli. |
| Specificity : | BLNK Monoclonal Antibody detects endogenous levels of BLNK protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Monoclonal, Mouse |
| Dilution : | WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. Flow cytometry: 1:200 - 1:400. ELISA: 1:10000. Not yet tested in other applications. |
| Purification : | Affinity purification |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |

Molecularweight : 50kD

Cell Pathway : B_Cell_Antigen;Primary immunodeficiency;

P References : 1. J Biol Chem. 2009 Apr 10;284(15):9804-13.
2. Cancer Sci. 2008 Dec;99(12):2444-54.


Background : This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012],

Function : disease:Defects in BLNK are the cause of hypoglobulinemia and absent B-cells [MIM:604515]. This is a developmental blockage at the pro- to pre-B-cell transition.,disease:In 6 of 34 childhood pre-B acute lymphoblastic leukemia (ALL) samples that were tested showed a complete loss or drastic reduction of BLNK expression.,function:Functions as a central linker protein that bridges kinases associated with the B-cell receptor (BCR) with a multitude of signaling pathways, regulating biological outcomes of B-cell function and development. Plays a role in the activation of ERK/EPHB2, MAP kinase p38 and JNK. Modulates AP1 activation. Important for the activation of NF-kappa-B and NFAT. Plays an important role in BCR-mediated PLCG1 and PLCG2 activation and Ca(2+) mobilization and is required for trafficking of the BCR to late endosomes. However, does not seem to be required for pre-BCR-mediated ac


Subcellular Location : Cytoplasm . Cell membrane . BCR activation results in the translocation to membrane fraction.

Expression : Expressed in B-cell lineage and fibroblast cell lines (at protein level). Highest levels of expression in the spleen, with lower levels in the liver, kidney, pancreas, small intestines and colon.


Products Images



Western Blot analysis using BLNK Monoclonal Antibody against NIH/3T3 (1) and BCBL-1 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human cervical cancer tissues with DAB staining using BLNK Monoclonal Antibody.



Immunofluorescence analysis of HepG2 cells using BLNK Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Flow cytometric analysis of NIH/3T3 cells using BLNK Monoclonal Antibody (green) and negative control (purple).

