

Protein Z Monoclonal Antibody

Catalog No: YM0538

Reactivity: Human

Applications: WB;ELISA

Target: Protein Z

Gene Name: PROZ

Protein Name: Vitamin K-dependent protein Z

P22891

Q9CQW3

Human Gene Id: 8858

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: Purified recombinant fragment of Protein Z expressed in E. Coli.

Specificity: Protein Z Monoclonal Antibody detects endogenous levels of Protein Z protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.

Purification : Affinity purification

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 45kD

P References: 1. J Thromb Haemost. 2005 Mar;3(3):497-501.

2. Blood Coagul Fibrinolysis. 2008 Jan;19(1):23-5.



Background:

This gene encodes a liver vitamin K-dependent glycoprotein that is synthesized in the liver and secreted into the plasma. The encoded protein plays a role in regulating blood coagulation by complexing with protein Z-dependent protease inhibitor to directly inhibit activated factor X at the phospholipid surface. Deficiencies in this protein are associated with an increased risk of ischemic arterial diseases and fetal loss. Mutations in this gene are the cause of protein Z deficiency. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jan 2012],

Function:

caution:Although homologous with the vitamin K-dependent clotting factors, it has lost two of the essential catalytic residues and has no enzymatic activity.,function:Appears to assist hemostasis by binding thrombin and promoting its association with phospholipid vesicles.,PTM:The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 Gla (gamma-carboxy-glutamate) domain.,similarity:Contains 1 peptidase S1 domain.,similarity:Contains 2 EGF-like domains.,tissue specificity:Plasma.,


Subcellular Location:

Secreted.

Expression:

Plasma.

Products Images



Western Blot analysis using Protein Z Monoclonal Antibody against human plasma (1).