

Tubulin β Monoclonal Antibody

Catalog No: YM1110

Reactivity: Human; Mouse; Rat; Bovine; Rabbit

Applications: WB;IP

Target: Tubulin β

Fields: >>Phagosome;>>Gap junction;>>Alzheimer disease;>>Parkinson

disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Pathogenic

Escherichia coli infection;>>Salmonella infection

Gene Name: TUBB1

Protein Name: Tubulin beta-1 chain

P07437

A2AQ07

Human Gene Id: 81027

Human Swiss Prot

No:

Mouse Gene Id: 545486

Mouse Swiss Prot

No:

Immunogen: Purified recombinant human Tubulin β protein fragments expressed in E.coli.

Specificity: Tubulin β Monoclonal Antibody detects endogenous levels of Tubulin β protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution: WB 1:1000 - 1:2000. Not yet tested in other applications.

Purification : Affinity purification

Concentration: 1 mg/ml

1/2



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 50kD

Cell Pathway : Protein_Acetylation

Background: This gene encodes a member of the beta tubulin protein family. Beta tubulins are

one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is specifically expressed in platelets and megakaryocytes and may be involved in proplatelet production and platelet release. A mutations in this gene is associated with autosomal dominant

macrothrombocytopenia. Two pseudogenes of this gene are found on

chromosome Y.[provided by RefSeq, Jul 2010],

Function: function: Tubulin is the major constituent of microtubules. It binds two moles of

GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,similarity:Belongs to the tubulin

family., subunit: Dimer of alpha and beta chains.,


Subcellular Location:

Cytoplasm, cytoskeleton.

Expression: Hematopoietic cell-specific. Major isotype in leukocytes, where it represents

50% of all beta-tubulins.

Products Images



Western Blot analysis using Tubulin β Monoclonal Antibody against A431, HeLa, Jurkat cell lysate.