

**SOX-2 Monoclonal Antibody**

<b>Catalog No :</b>	YM0594
<b>Reactivity :</b>	Human
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	SOX-2
<b>Fields :</b>	>>Hippo signaling pathway;>>Signaling pathways regulating pluripotency of stem cells
<b>Gene Name :</b>	SOX2
<b>Protein Name :</b>	Transcription factor SOX-2
<b>Human Gene Id :</b>	6657
<b>Human Swiss Prot No :</b>	P48431
<b>Mouse Swiss Prot No :</b>	P48432
<b>Immunogen :</b>	Purified recombinant fragment of human SOX-2 expressed in E. Coli.
<b>Specificity :</b>	SOX-2 Monoclonal Antibody detects endogenous levels of SOX-2 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Monoclonal, Mouse
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	Affinity purification
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	34kD

**P References :**

1. Proc Natl Acad Sci U S A. 2008 Nov 25;105(47):18396-401.
  2. J Biol Chem. 2008 Nov 28;283(48):33730-5.
  3. Nature. 2008 Oct 23;455(7216):1124-8.
- 

**Background :**

SRY-box 2(SOX2) Homo sapiens This intronless gene encodes a member of the SRY-related HMG-box (SOX) family of transcription factors involved in the regulation of embryonic development and in the determination of cell fate. The product of this gene is required for stem-cell maintenance in the central nervous system, and also regulates gene expression in the stomach. Mutations in this gene have been associated with optic nerve hypoplasia and with syndromic microphthalmia, a severe form of structural eye malformation. This gene lies within an intron of another gene called SOX2 overlapping transcript (SOX2OT). [provided by RefSeq, Jul 2008],

---

**Function :**

disease:Defects in SOX2 are the cause of microphthalmia syndromic type 3 (MCOPS3) [MIM:206900]. Microphthalmia is a clinically heterogeneous disorder of eye formation, ranging from small size of a single eye to complete bilateral absence of ocular tissues (anophthalmia). In many cases, microphthalmia/anophthalmia occurs in association with syndromes that include non-ocular abnormalities. MCOPS3 is characterized by the rare association of malformations including uni- or bilateral anophthalmia or microphthalmia, and esophageal atresia with trachoesophageal fistula.,function:Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.,online information:Sox2 entry,PTM:Sumoylation inhibits bin

---

**Subcellular Location :**

Nucleus speckle . Cytoplasm . Nucleus . Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (By similarity). Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH) (By similarity). Colocalizes with SOX6 in speckles. Colocalizes with CAML in the nucleus (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (By similarity). .


---

**Expression :**


Fetal brain,Lung,Retina,

---


## Products Images



Western Blot analysis using SOX-2 Monoclonal Antibody against HEK293 (1) and SOX2-hlgGfc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded lung cancer tissues (left) and esophageal cancer tissues (right) with DAB staining using SOX-2 Monoclonal Antibody.



Immunofluorescence analysis of NTERA-2 cells using SOX-2 Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.