

TORC3 Monoclonal Antibody

Catalog No :	YM0626
Reactivity :	Human;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	TORC3
Fields :	>>Human T-cell leukemia virus 1 infection
Gene Name :	CRTC3
Protein Name :	CREB-regulated transcription coactivator 3
Human Gene Id :	64784
Human Swiss Prot No :	Q6UUV7
Mouse Swiss Prot No :	Q91X84
Immunogen :	Purified recombinant fragment of human TORC3 expressed in E. Coli.
Specificity :	TORC3 Monoclonal Antibody detects endogenous levels of TORC3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Monoclonal, Mouse
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	Affinity purification
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	67kD

P References :

1. Mod Pathol. 2009 Dec;22(12):1575-81.
 2. Genes Chromosomes Cancer. 2008 Mar;47(3):203-6.
-

Background :

This gene is a member of the CREB regulated transcription coactivator gene family. This family regulates CREB-dependent gene transcription in a phosphorylation-independent manner and may be selective for cAMP-responsive genes. The protein encoded by this gene may induce mitochondrial biogenesis and attenuate catecholamine signaling in adipose tissue. A translocation event between this gene and Notch coactivator mastermind-like gene 2, which results in a fusion protein, has been reported in mucoepidermoid carcinomas. Alternative splicing results in multiple transcript variants that encode different protein isoforms. [provided by RefSeq, Jul 2012],

Function :

function:Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).,similarity:Belongs to the TORC family.,subcellular location:Appears to be mainly nuclear.,subunit:Binding, as a tetramer, through its N-terminal region, with the bZIP domain of CREB1 enhances recruitment of TAF4 to the promoter. 'Arg-314' in the bZIP domai


Subcellular Location :

Nucleus . Cytoplasm . Appears to be mainly nuclear (PubMed:15454081). Translocates to the nucleus following adenylyl cyclase or MAP kinase activation (PubMed:30611118). .


Expression :

Predominantly expressed in B and T lymphocytes. Highest levels in lung. Also expressed in brain, colon, heart, kidney, ovary, and prostate. Weak expression in liver, pancreas, muscle, small intestine, spleen and stomach.


Products Images



Western Blot analysis using TORC3 Monoclonal Antibody against HeLa (1), Jurkat (2), Cos7 (3) and MCF-7 (4) cell lysate.



Immunohistochemistry analysis of paraffin-embedded breast cancer (left) and ovarian cancer (right) with DAB staining using TORC3 Monoclonal Antibody.



Immunofluorescence analysis of NTERA-2 cells using TORC3 Monoclonal Antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

