

Bcl-XL (PT0244R) PT® Rabbit mAb

Catalog No: YM8153

Reactivity: Human; Mouse; Rat;

Applications: WB;IHC;IF;IP;ELISA

Target: Bcl-x

Fields: >>EGFR tyrosine kinase inhibitor resistance;>>Platinum drug resistance;>>Ras

signaling pathway;>>NF-kappa B signaling pathway;>>p53 signaling

pathway;>>Mitophagy - animal;>>Autophagy - animal;>>PI3K-Akt signaling pathway;>>Apoptosis;>>Apoptosis - multiple species;>>NOD-like receptor

signaling pathway;>>JAK-STAT signaling pathway;>>Parkinson

disease;>>Amyotrophic lateral sclerosis;>>Pathways of neurodegeneration - multiple diseases;>>Shigellosis;>>Toxoplasmosis;>>Measles;>>Human T-cell

leukemia virus 1 infection;>>Herpes simplex virus 1 infection;>>Human immunodeficiency virus 1 infection;>>Pathways in cancer;>>Transcriptional

misregulation in cancer;>>Pancreatic cancer;>>Chronic myeloid

leukemia;>>Small cell lung cancer;>>Hepatocellular carcinoma;>>Lipid and

atherosclerosis

Q64373

Gene Name: BCL2L1

Protein Name: Bcl-2-like protein 1

Human Gene Id: 598

Human Swiss Prot Q07817

No:

Mouse Gene Id: 12048

Mouse Swiss Prot

No:

Rat Gene Id: 24888

Rat Swiss Prot No: P53563

Specificity: endogenous

1/3



Formulation: PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Source: Monoclonal, rabbit, IgG, Kappa

Dilution: IHC 1:200-1:1000,WB 1:1000-1:5000,IF 1:200-1:1000,ELISA

1:5000-1:20000, IP 1:50-1:200,

Purification: Protein A

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 26kD

Observed Band: 30kD

Cell Pathway: Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;Jak_STAT;

Amyotrophic lateral sclerosis (ALS);Pathways in cancer;Pancreatic

cancer; Chronic myeloid leukemia; Small cell lung cancer;

Background: The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2

family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening.

VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by

mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an

apoptotic activator. [provided by RefSeq, Dec 2015],

Function: domain: The BH4 motif is required for anti-apoptotic activity. The BH1 and BH2

motifs are required for both heterodimerization with other Bcl-2 family members and for repression of cell death., function: Potent inhibitor of cell death. Isoform Bcl-X(L) anti-apoptotic activity is inhibited by association with SIVA isoform 1. Inhibits activation of caspases (By similarity). Appears to regulate cell death by blocking the voltage-dependent anion channnel (VDAC) by binding to it and preventing the

release of the caspase activator, cytochrome c, from the mitochondrial membrane. The Bcl-X(S) isoform promotes apoptosis.,PTM:Proteolytically cleaved by caspases during apoptosis. The cleaved protein, lacking the BH4 motif, has pro-apoptotic activity.,similarity:Belongs to the Bcl-2 family.,subcellular location:Mitochondrial membranes and perinuclear envelope.,subunit:Bcl-X(L)

forms homodimers, and het

Subcellular Location:

Cytoplasm


2/3




Expression:

Bcl-X(S) is expressed at high levels in cells that undergo a high rate of turnover, such as developing lymphocytes. In contrast, Bcl-X(L) is found in tissues containing long-lived postmitotic cells, such as adult brain.


Products Images



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Bcl-XL (PT0244R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1:3T3-L1 Lane 2: C6 Lane 3: HeLa Lane 4: K562 Predicted band size: 26kDa Observed band size: 30kDa



Human colon carcinoma was stained with anti-Bcl-XL (PT0244R) rabbit antibody



Human colon carcinoma was stained with anti-Bcl-XL (PT0244R) rabbit antibody